



# Influence of Different Levels of Hydrogel on Root Nodulation in Chickpea (*Cicer arietinum* L.)

Shalini Singh <sup>a\*</sup> and Richa Sharma <sup>a</sup>

<sup>a</sup> Department of Biological Sciences, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, India.

## Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i94956>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122204>

Original Research Article

Received: 19/06/2024

Accepted: 21/08/2024

Published: 28/08/2024

## ABSTRACT

This study investigated the impact of varying levels of hydrogel on the root nodules of Chickpea (*Cicer arietinum* L.) during Rabi seasons of 2019-20 and 2021-22 at the Department of Biological Sciences, Sam Higginbottom Institute of Agriculture, Technology, and Sciences, (U.P). The experimental design was a randomized block design with four levels of hydrogel (25%, 50%, 75% and 100%) and two levels of irrigation (one irrigation and two irrigation) and all treatments replicated thrice. Observations on the number of nodules per plant, active nodules, and inactive nodules were recorded at 30, 60, and 90 days after sowing (DAS) and at maturity and analysed using Fisher's ANOVA. The results indicated that hydrogel application significantly affected nodule formation at all growth stages across both years. The highest number of nodules, active nodules, and inactive nodules were recorded in treatment T<sub>10</sub> (2 irrigations + 100% hydrogel), followed by T<sub>9</sub> (2 irrigations + 75% hydrogel), in both years. The enhanced nodule formation with hydrogel application was

\*Corresponding author: E-mail: shalinisingh2909@gmail.com;

attributed to improved water availability to the roots during periods of water scarcity, highlighting hydrogel's role in optimizing nitrogen availability and promoting chickpea growth, which is crucial for enhancing soil fertility.

**Keywords:** Root nodulation chickpea; agricultural productivity; rain-fed crops; leguminous plant.

## 1. INTRODUCTION

Being a tropical country, India has seen a variety of changes in weather patterns and a rise in temperature resulted of change in climate and global warming. Increased transpiration caused by these changes reduces soil water levels, which has an impact on agricultural productivity. Water is one of necessary material for a plant to develop and advance [1]. In agriculture, irrigation and rainfall are the two primary water sources. 65% of the food produced globally comes from crops that are grown using irrigation, while the remaining 35% comes from rain-fed crops. Irrigation covers just 17% of all planted land. (Rosegrant, 2002 and Hanjra, 2010). As a result, the majority of land in agriculture is dependent on natural precipitation. Thus, an unprecedented drought is occurring in many of the world's food producing regions as a resulted of variation in the global rain pattern and a rise in local temperature. (FAO, 2008). Chickpeas possess deeply penetrating roots that enable them to access and utilize the limited moisture available in the soil, making them valuable for improving soil fertility [2]. However, the challenges posed by salinized soils underscore the need for developing more resilient rhizobial inoculants that can thrive in such conditions, thereby enhancing the efficiency of nitrogen fixation and ultimately contributing to sustainable agriculture in saline environments.

Nitrogen fixation of leguminous plant is an important aspect of sustainable agriculture by which the atmospheric  $N_2$  is converted to ammonia with the aid of a key enzyme called nitrogenase [3,4]. It is achieved by bacteria inside the cells of *de novo* formed organs, the nodules, which usually develop on roots of various leguminous plants. This process is resulted from the complex interaction between the host plant and rhizobia. The chickpea, like other legumes, is important for environmentally friendly and sustainable farming [5,6,7]. It may form a symbiotic relationship with rhizobia that can convert atmospheric nitrogen into ammonia that plants can directly absorb, increasing the productivity of this crop and succeeding cereal crops and enhancing soil fertility by fixing

nitrogen at rates of up to 140 kg/ha/year [8], decreases the need of chemical fertilisers greatly, which lowers the amount of greenhouse gas emissions and water pollution per hectare each year [9]. By boosting the synthesis of phytohormones and mineral absorption, several rhizobial strains have been shown to be able to stimulate plant growth and development [10]. Chickpea is regarded as a host that restricts nodulation (Rivas *et al.*, 2007) [11,12]. The availability of nitrogen in agriculture has a significant impact on both output and product quality. In the natural world, plants either assimilate nitrate and ammonium or get nitrogen from di-nitrogen by collaborating with bacteria that fix nitrogen. The most significant symbiosis in terms of biological nitrogen fixation is between legumes and rhizobia, which together produce roughly 200 million tonnes of nitrogen annually [13,14,15,16]. The bacterial microsymbionts enter individual plant cells and settle in compartments encircled by a plant membrane to form a symbiosis necessary for nodulation [17]. Although the symbiotic nitrogen (N) fixation between legumes and Rhizobium is a crucial biological characteristic and the cornerstone of increasing soil fertility, there are currently few efficient rhizobial inoculants that can adapt to salinized soils. Previous research has demonstrated that drought and salt stress significantly reduced plant biomass accumulation (root and shoot), nodule growth, nitrogenase activity, and chickpea production [18,19]. Due to their deeply penetrating roots, which allow them to make use of the scarce moisture that is available; chickpeas can improve the fertility of the soil [2].

## 2. MATERIALS AND METHODS

The field experiment of present investigation entitled "Impact of different levels of hydrogel on root nodules of Chickpea (*Cicer arietinum* L.)" was conducted during "rabi season" 2019-20 and 2021-22 under Department of Biological Sciences, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad (U.P). The climate of the experimental site is characterised by semi-arid type with hot and dry summer from April to June, hot and

humid from July to September and cold winter from November to January. The experiment was conducted by the randomised block design with combinations of four level of hydrogel and two level of irrigation i.e. T<sub>0</sub> (3 irrigation), T<sub>1</sub> (1 irrigatin) T<sub>2</sub> (1 irrigation + 25% hydrogel) T<sub>3</sub> (1 irrigation + 50% hydrogel), T<sub>4</sub> (1 irrigation + 75% hydrogel), T<sub>5</sub> (1 irrigation + 100% hydrogel), T<sub>6</sub> (2 irrigatin) T<sub>7</sub> (2 irrigation + 25% hydrogel), T<sub>8</sub> (2 irrigation + 50% hydrogel), T<sub>9</sub> (2 irrigation + 75% hydrogel), T<sub>10</sub> (2 irrigation + 100% hydrogel) are replicated thrice. During the crop growth period, the following observations were recorded time to time. Number of nodules/plants: After digging out three plants from randomly in each plot, the number of nodules were counted at regular intervals of 30, 60, 90 DAS and at maturity. Active nodules: Active nodules were counted on the basis of colour of the nodules. Inactive nodules: Inactive nodules were counted by subtracting Active nodules from Total number of nodules. Fisher's ANOVA technique and least significance difference (LSD) test at 5% probability level were used to compare differences among treatment means.

### 3. RESULTS AND DISCUSSION


Application of Hydrogel to soil for cultivation of many crops under water scarce areas is proven beneficial. Many parts of studied area have water scarcity, hence present experimentation was conducted to assess the growth and yield performance of Chickpea (*Cicer arietinum* L.). The result has been presented and graphically illustrated using Table 1 and Fig. 1. The effect of application of hydrogel was found to be significant at 30DAS, 60 DAS, 90 DAS and at

maturity in both the years. The maximum no. of Nodules 24.82, 32.37, 25.0 and 24.5 were recorded in treatment T<sub>10</sub> in the year 2020, 2021and in pooled respectively and followed by treatment combination T<sub>9</sub> which is at par and recorded No. of Nodules in both the years respectively.

The effect of application of hydrogel on No. of Active Nodules was found to be significant at 30DAS, 60 DAS, 90 DAS and at maturity in both the years. The maximum No. of Active Nodules 22.52, 30:20.13:19 was recorded in treatment T<sub>10</sub> in the year 2020, 2021and in pooled respectively and followed by treatment combination T<sub>9</sub> which is at par and recorded No. of Active Nodules in both the years respectively.

The effect of application of hydrogel on No. of Inactive Nodules was found significant at 30DAS, 60 DAS, 90 DAS and at maturity in both the years. The maximum No. of Inactive Nodules 3.52, 5.23, 9.17 and 7.82 was recorded in treatment T<sub>10</sub> in the year 2020, 2021and in pooled respectively and followed by treatment combination T<sub>9</sub> which is at par and recorded No. of Inactive Nodules in both the years respectively.

This may due to the availability of water to the roots through hydrogel during water scarcity to the plants. The availability of nitrogen in agriculture has a significant impact on both output and product quality [13,14]. Chickpeas possess deeply penetrating roots that enable them to access and utilize the limited moisture available in the soil, making them valuable for improving soil fertility [2].



**Fig. 1. Influence of different levels of hydrogel on root nodules of Chickpea (*Cicer arietinum* L.)**

**Table 1. Influence of different levels of hydrogel on root nodules of Chickpea (*Cicer arietinum* L.)**

Treatments	No. of nodules (Nos)				No. of active nodules (Nos)				No. of inactive nodules (Nos)			
	30DAS	60DAS	90DAS	At harvest	30DAS	60DAS	90DAS	At harvest	30DAS	60DAS	90DAS	At harvest
T <sub>0</sub>	21.12	26.23	23.13	21.77	19.42	24.37	17.35	14.9	1.7	1.85	5.77	6.82
T <sub>1</sub>	13.6	18.27	14.83	14.37	10.08	16.4	7.45	6.67	3.52	1.88	7.38	7.68
T <sub>2</sub>	13.47	19.4	14.3	14	10.92	18.5	8.1	7.27	2.55	0.93	6.2	6.72
T <sub>3</sub>	13.73	20.2	14.9	14.5	11.45	18.63	8.33	7.67	2.28	1.58	6.55	6.82
T <sub>4</sub>	13.2	20.4	14.5	13.8	11.72	18.13	8.7	8.13	1.48	2.28	5.8	5.62
T <sub>5</sub>	14.13	20.73	15.17	14.63	12.18	17.2	9.2	8.7	1.95	3.53	5.97	5.92
T <sub>6</sub>	14.93	24.33	16.47	16.2	12.83	19.13	9.23	8.35	2.1	5.23	7.23	7.82
T <sub>7</sub>	20.93	24.3	19.37	19.7	17.45	20.2	10.2	9.57	3.48	4.12	9.17	10.1
T <sub>8</sub>	24.18	30.33	24.13	23	20.8	27.4	18.23	17.3	3.38	2.95	5.88	5.7
T <sub>9</sub>	23.25	31.03	24.53	23.7	21.4	28.93	19.18	17.52	1.85	2.1	5.32	6.17
T <sub>10</sub>	24.82	32.37	25.0	24.5	22.52	30.0	20.13	19.0	2.3	2.37	4.87	5.47
C.D.	0.91	2.45	1.91	2	0.43	1.77	0.19	0.93	0.97	1.49	1.98	2.2
SE(m)	0.31	0.82	0.64	0.67	0.15	0.6	0.06	0.31	0.33	0.5	0.67	0.74
SE(d)	0.44	1.17	0.91	0.95	0.21	0.84	0.09	0.44	0.46	0.71	0.94	1.05
C.V.	2.97	5.87	5.94	6.42	1.62	4.75	0.88	4.74	23.3	33.07	18.14	18.89

#### 4. CONCLUSION

The results of present study concluded that hydrogel application significantly affected nodule formation at all growth stages across both years. The highest number of nodules, active nodules, and inactive nodules were recorded in treatment T<sub>10</sub> (2 irrigations + 100% hydrogel), followed by T<sub>9</sub> (2 irrigations + 75% hydrogel), in both years. The enhanced nodule formation with hydrogel application was attributed to improved water availability to the roots during periods of water scarcity, highlighting hydrogel's role in optimizing nitrogen availability and promoting chickpea growth, which is crucial for enhancing soil fertility.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Rijsberman FR. Water for food: corruption in irrigation systems. Global Corruption Report 2008. 2008;67.
2. Tripathi LK, Thomas T, Singh VJ, Gampala S, Kumar R. Effect of nitrogen and phosphorous application on soil nutrient balance in Chickpea (*Cicer arietinum* L) cultivation. Green Frming. 2015;6(2):319-322.
3. Udvardi M, Poole PS. Transport and metabolism in legume-rhizobia symbioses. Annu. Rev. Plant Biol. 2013;64:781–805.
4. Oldroyd GE, Murray JD, Poole PS, Downie JA. The rules of engagement in the legume-rhizobial symbiosis. Annu. Rev. Genet. 2011;45:119–144.
5. Erdemci İ, Aktaş H, Nadeem MA. Effect of fertilization and seed size on nodulation, yield and yield components of Chickpea (*Cicer Arietinum* L.). Applied Ecology and environmental Research; 2017.
6. Kumar J, Kumar S, Prakash V. Effect of biofertilizers and phosphorus levels on soil fertility, yield and nodulation in chickpea (*Cicer arietinum* L.). Journal of the Indian Society of Soil Science. 2019;67(2):199-203.
7. Kumar S, Tripathi DK, Bharose R, Kumar M, Kumar R. Effect of different fertility level and micronutrients on nodulation and nutrient uptake by chickpea. An Asian Journal of Soil Sci. 2016;11(1):62-66.
8. Flowers TJ, Gaur PM, Gowda CLL, Krishnamurthy, Samineni S, Siddique KM, et al. Salt sensitivity in Chickpea. Plant, Cell Environ. 2010;33:490-509.
9. Coba de la Pena T, Fedorova E, Pueyo JJ, Lucas MM. The symbiosome: legume and rhizobia co-evolution toward a nitrogen-fixing organelle?. Frontiers in plant science. 2018 Jan 22;8:2229.
10. Karthik S, Siva P, Balu KS, Suriyaprabha R, Rajendran V, Maaza M. Acalypha indica-mediated green synthesis of ZnO nanostructures under differential thermal treatment: Effect on textile coating, hydrophobicity, UV resistance, and antibacterial activity. Advanced Powder Technology. 2017 Dec 1;28(12):3184-94.
11. Laranjo M, Alexandre A, Rivas R, Velázquez E, Young JPW, Oliveira S. Chickpea rhizobia symbiosis genes are highly conserved across multiple *Mesorhizobium species*. FEMS Microbiol Ecol. 2008;66(2):391-400.
12. Suneja P, Dudeja SS, Dahiya P. Deciphering the phylogenetic relationships among rhizobia nodulating chickpea: A review. J. Appl. Biol. Biotechnol. 2016 May;4:61-70.
13. Graham PH, Vance CP. Legume importance and constraints to greater use. Plant Physiology. 2003;131:872-877.
14. Peoples MB, Brockwell J, Herridge DF, Rochester IJ, Alves BJR, Urquiaga S, et al. The contributions of nitrogen-fixing crop legumes to the productivity of agricultural systems. Symbiosis. 2009;48:1-17.
15. Gebremariam M, Tesfay T. Effect of P application rate and rhizobium inoculation on nodulation, growth, and yield performance of chickpea (*Cicer arietinum* L.). International Journal of Agronomy. 2021(1):8845489.
16. Singh G, Sekhon HS, Sharma P. Effect of irrigation and biofertilizer on water use, nodulation, growth and yield of chickpea (*Cicer arietinum* L.). Archives of Agronomy and Soil Science. 2011;57(7):715-726.
17. Jens Stougaard. Regulators and regulation of legume root nodule development. Plant Physiology. 2000;124:531-540.

18. Garg N, Baher N. Role of arbuscular mycorrhizal symbiosis in proline biosynthesis and metabolism of *Cicer arietinum* L.(chickpea) genotypes under salt stress. *Journal of plant growth regulation*. 2013 Dec;32:767-78.
19. Egamberdieva D, Lugtenberg B. Use of plant growth-promoting rhizobacteria to alleviate salinity stress in plants. *Use of Microbes for the Alleviation of Soil Stresses, Volume 1*. 2014:73-96.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/122204>