



Correlation Between Predictors of Obstructive Sleep Apnoea

Bright Otohile^{1*}, Odiase Oboite² and Okubokekeme Otoru Otoru³

¹Department of Otorhinolaryngology, Federal Medical Centre, Yenagoa, Nigeria.

²Department of Anaesthesia, Federal Medical Centre, Yenagoa, Nigeria.

³Department of Family Medicine, Federal Medical Centre, Yenagoa, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author BO designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors OO and OOO managed the analyses of the study. Author OOO managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2018/46651

Editor(s):

(1) Dr. Andrea S. Melani, Department of Cardiothoracic Disease, Azienda Ospedaliera Universitaria Senese, Italy.

Reviewers:

(1) Serghei Covantev, State University of Medicine and Pharmacy "Nicole Testemitanu", Republic of Moldova.

(2) Sidika Deniz Yalim, Adana City Training and Research Hospital, Turkey.
Complete Peer review History: <http://www.sdiarticle3.com/review-history/46651>

Original Research Article

Received 28 October 2018

Accepted 13 January 2019

Published 29 January 2019

ABSTRACT

Background: Obstructive sleep apnoea is becoming a respiratory disorder of public health concern. Different obesity indices such as Mallampati score, Body Mass Index, neck circumference have been identified in the literature as predictors of obstructive sleep apnoea. Studies on the relationship between these variables is however limited particularly in Nigeria. The aim of this study was to evaluate the correlation between Mallampati score, Body Mass Index and neck circumference which are predictors of obstructive sleep apnoea.

Methods: The study was done among commercial drivers. The risk of obstructive sleep apnoea was assessed with Berlin questionnaire. The participants were assessed for Mallampati score, Body Mass Index; their neck circumference was measured and the correlation between these variables was done. Data collated was analyzed with statistical package for social sciences (SPSS) Version 20.0 software.

Results: There were 105 drivers recruited into this study, they were all males with a mean age of 44.8 ± 12.03 years and 50.5% of the drivers had high risk of sleep apnoea. There was a positive

*Corresponding author: E-mail: otoghilebright@gmail.com;

correlation between Mallampati score, Body Mass Index and neck circumference.

Conclusion: Mallampati score increase with increase in Body Mass Index and neck circumference which are all predictors of obstructive sleep apnoea.

Keywords: Correlation; Mallampati; BMI; neck circumference; sleep.

1. INTRODUCTION

Obstructive sleep apnoea is a common respiratory disorder. Although the pathophysiology of obstructive sleep apnoea is yet to be clearly understood, there are anatomical abnormalities that could possibly result to upper airway obstruction during sleep. The onset of inspiration has been demonstrated to trigger a reflex increase in the electromyographic (EMG) activity of the pharyngeal muscles [1]. These muscles are activated during daytime respiration and their normal tone keeps the airway open thereby preventing collapse of the upper airway; but during REM sleep there is hypotonia of these muscles. In patients with obstructive apnoea, the EMG activity of the pharyngeal muscles decreases further during obstructive episode [2]. It is essential to note that any reduction in tone of the upper airway muscles or any incoordination between reflex EMG activity in these muscles and onset of inspiration causes an increase susceptibility of the upper airway to collapse. However, apart from these aforementioned factors, risk factors for obstructive sleep apnoea include increasing age, sex, alcohol, hypertension, diabetes, neck circumference, Mallampati and Body Mass Index (BMI) [3-5]. These are the predictors of obstructive sleep apnoea. Of these factors, neck circumference, Mallampati and Body Mass Index (BMI) are obesity indices [6-7]. Screening for Obstructive sleep apnoea in drivers is very important as undiagnosed cases among them poses risk for the society due to their job. Several studies in Nigeria have shown that commercial car drivers are particularly at risk of Obstructive sleep apnoea [8-10]. However, these studies did not consider the relationship between these obesity indices. Hence, the aim of this study was to examine the correlation between neck circumference, Mallampati and Body Mass Index which are all predictors of Obstructive sleep apnoea and using commercial car drivers as case study.

2. METHODS

2.1 Study Design

This is a prospective, descriptive study involving commercial car drivers.

2.2 Study Setting

Study was done in Ile-Ife in Osun state in the south western part of Nigeria.

2.3 Study Protocol

All commercial drivers in Ile-Ife who gave consent were recruited into the study. The Berlin questionnaire was administered to all the participants and this was used to assess the risk of obstructive sleep apnoea. The Berlin questionnaire contains 10 questions which are scored positive or negative and the questions are divided into 3 categories. A participant was said to have high risk of sleep apnoea if the score is positive in 2 or more categories of questions and low risk if the score is positive in one or no category of questions. The neck circumference, Mallampati and BMI of all participants were assessed and recorded. The mouth of the participants were inspected while widely opened with tongue fully extruded and sitting upright, and Mallampati classification of what is visible was done as follows:

- Class I – soft palate, anterior and posterior faucial pillars and uvula.
- Class II – faucial pillars, soft palate and uvula.
- Class III – soft palate and base of uvula.
- Class IV – soft palate not visible.

Their weight, height and blood pressure were measured using a weighing scale, Stadiometer and sphygmomanometer respectively. The Body Mass Index (BMI) was calculated and graded as underweight if less than 18.5, normal from 18.5 - 24.9, overweight from 25.0 - 29.9, obese from 30.0 - 39.9 and from 40 and above was regarded as morbidly obese. Their neck circumference was measured in centimetres using a metric tape at the level of the thyrohyoid membrane. Neck circumference greater than 43.35 cm (17 inches) was regarded as abnormal.

2.4 Ethical Approval

Ethical Clearance was obtained from the Ethics and Research Committee of the Obafemi

Awolowo University Teaching Hospitals Complex, Ile-Ife.

2.5 Data Management

Data collected was analyzed using Statistical Package for Social Sciences version 20.0 and a 'p' value of less than 0.05 was accepted as statistically significant. Results were subsequently presented in descriptive format.

3. RESULTS

One hundred and five commercial car drivers were included in the study. They were all males and the mean age ($\bar{X} \pm SD$) was 44.8±12.03 years. Average Neck circumference (cm) and BMI (Kg/m²) were 39.0±3.3 and 25.4±4.6 respectively. A total of 53(50.5%) drivers had high risk of sleep apnoea while 52(49.5%) of them had low risk. Table 1 show the correlation between neck circumference, Mallampati and BMI and it reveals that Mallampati increases with increase in neck circumference and BMI. Table 2 shows that as BMI increases there is higher

percentage of participants with neck circumference above 43.35 cm and Table 3 shows that as Mallampati score increases, there are more proportion of participants with neck circumference above 43.35 cm.

4. DISCUSSION

Findings from this study showed that a significant proportion of the participants had high risk of sleep apnoea. Of all the several predisposing factors to obstructive sleep apnoea, Mallampati, neck circumference and BMI have been shown to contribute more when compared to others [8]. This suggests that there is a relationship between the body anthropometry and obstructive sleep apnoea [11,12]. Increasing neck circumference, BMI and Mallampati is associated with a potential crowding of the oropharynx which consequently result in a narrowing of the air column which in conjunction with hypotonia of the pharyngeal muscles during sleep, could cause obstructive sleep apnoea.

Table 1. Showing the correlations between Mallampati, BMI and Neck Circumference

Variable	Mallampati	BMI	Neck circumference
Mallampati	1		
BMI	0.594**	1	
Neck circumference	0.596**	0.792**	1

** Correlation is significant at the 0.01 level (2-tailed)

Table 2. Association between neck circumference and BMI

BMI	Neck circumference		Chi-square	P-value
	≤43.35 cm (n, %)	>43.35 cm (n, %)		
<18.5	5 (100)	0	6.200	0.001
18.5 -24.9	48 (98.0)	1(2.0)		
25.0-29.9	32 (97.0)	1(3.0)		
30.0-39.9	11 (61.1)	7(38.9)		

Table 3. Association between neck circumference and Mallampati score

Mallampati score	Neck circumference		Chi-square	P-value
	≤43.35 cm (n, %)	>43.35 cm (n, %)		
I	49(98.0)	1(2.0)	39.000	0.001
II	32(97)	1(3)		
III	14(66.7)	6(33.3)		
IV	1(50)	1(50)		

Results from this study shows that there is a positive correlation between Mallampati, BMI and neck circumference. This shows that even as these obesity indices such as Mallampati, BMI, and neck circumference predict obstructive sleep apnoea, they also predict and interrelate with each other as well. There is paucity of comparable local studies. However, community based study from India shows that neck circumference is a predictor of BMI [13]. Kumar et al. in another study in India also corroborate these findings [14]. Studies from Iran and United States of America also had similar results [15,16]. Table 2 from this study showed that there was an increase in the percentage of participants with neck circumference greater than 43.35cm as BMI increases. Consequently, neck circumference being an indicator of upper body fat distribution can be used as measure obesity. Neck circumference has a positive correlation with Mallampati scores as found in this study. Table 3 show that the percentage of participants with neck circumference above 17 inches increases with higher Mallampati scores. Higher Mallampati scores indicate crowding of the oropharynx which has been seen to be related to neck circumference and other obesity indices. This probably explains the positive correlation between these variables demonstrated in this study. Menon et al. also demonstrated that neck circumference has positive correlation with Body Mass Index and Mallampati [7]. Apart from neck circumference having a positive correlation with Body Mass Index and Mallampati, both Body Mass Index and Mallampati also correlate with each other positively as shown in Table 1. The same study by Menon et al. showed that Body Mass Index has positive correlation with Mallampati scores [7]. This shows that neck circumference, BMI and Mallampati all correlate with each other and can predict each other even as they are also predictors of obstructive sleep apnoea. Hence, each of these variables can serve as a guide when assessing obesity, obstructive sleep apnoea. They can also serve when assessing the airway for possibility of difficulty in endotracheal intubation. Though Mallampati scoring may require some level of clinical training, neck circumference and Body Mass Index can easily be measured and calculated respectively by anyone who is conversant with figures and measurements. Any of these variables could be measured and used to predict obstructive sleep apnoea among commercial car drivers as used in this study. Studies have shown that these variables are risk factors for obstructive sleep apnoea among

drivers in this study location [8,9]. However, the correlation between other possible predictors of obstructive sleep apnoea such as age, sex, blood glucose, blood pressure were not considered in this study. Polysomnography is the gold standard for the diagnosis of obstructive sleep apnoea [17]. This was also not done in this study. Indeed, nevertheless polysomnography is costly and now obstructive sleep apnoea is often diagnosed with cheaper methods [18]. Questionnaires are becoming more widely used [19]. However, this study considered the correlation between some of the possible predictors of obstructive sleep apnoea which to the best of the authors' knowledge was not done in other previous similar local studies.

5. CONCLUSION

This study found that there are positive correlations between Mallampati, BMI and neck circumference. Therefore, any of these indices could be measured and easily be used as a predictor of obstructive sleep apnoea. We recommend study of the correlations between Mallampati, BMI, neck circumference and other predictors of obstructive sleep apnoea such as blood sugar, blood pressure, sex.

CONSENT

Consent was obtained from all participants.

ETHICAL APPROVAL

The study was done in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mathur R, Jan MA, Mastropasquia B, Douglas NJ. Upper airway dilator muscle reflex to negative pressure. British Sleep Society Meeting. University of Leicester. 1992;29.
2. Guilleminault C, Hill MW, Simmons FB, Dement WC. Obstructive sleep apnea: Electromyographic and fiberoptic studies. *Exp Neurol.* 1978;62(1):48-67.

3. Yusoff MF, Baki MM, Mohamed N, Mohamed AS, Yunus MR, Ami M, et al. Obstructive sleep apnoea among express bus drivers in Malaysia: Important indicators for screening. *Traffic Inj Prev.* 2010;11(6):594-599.
4. Sogebi OA, Oyewole EA, Soga-Peters OO. Sleep disordered breathing (SDB) experiences associated with snoring among adult Nigerians. *Afr Health Sci.* 2011;11(3):309-314.
5. Liistra D, Rombaux Ph, Belge C, Dury M, Aubert G, Rodenstein DO. High Mallampati score and nasal obstruction are associated risk factors for obstructive sleep apnoea. *Eur Respir J.* 2003;21:248-252.
6. Saka M, Türker P, Ercan A, Kiziltan G, Baş M. Is neck circumference measurement an indicator for abdominal obesity? A pilot study on Turkish Adults. *Afr Health Sci.* 2014;14(3):570-575.
7. Menon SM, Sampangiramaiah S, Mathew M. Cross sectional observational study performed to see for relation of Mallampati score and extended Mallampati score with body mass index. *Journal of Clinical and Diagnostic Research.* 2017;11(5):1-3.
8. Otoghile B, Amusa YB, Lasisi AO, Kuni JI, Otoru OO. Risk of obstructive sleep apnoea among commercial car drivers in a Nigerian City. *Nig Del Med J.* 2018;2(2): 42-48.
9. Obaseki DO, Erhabor GE, Obaseki JE, Abidoye I, Adebisi A, Olaiya B. Obstructive sleep apnea, excessive daytime sleepiness, and road traffic accidents among interstate commercial vehicle drivers in Nigeria. *J Resp Med;* 2014. DOI:<http://dx.doi/10.1155/2014/580264>
10. Ozoh OB, Okubadejo NU, Akanbi MO, Dania MG. High-risk of obstructive sleep apnoea and excessive daytime sleepiness among commercial intra-city drivers in Lagos metropolis. *Niger Med J.* 2013; 54(4):224-229.
11. Yildirim Y, Yilmaz S, Güven M, Kılınc F, Ali VK, Yilmaz Z, et al. Evaluation of anthropometric and metabolic parameters in obstructive sleep apnea. *Pulmonary Medicine.* 2015;1-6.
12. Kim SE, Park BO, Park SH, Shin KJ, Ha SY, Park J, Park KM. Predictors for presence and severity of obstructive sleep apnea in snoring patients and significance of neck circumference. *J Sleep Med.* 2015; 12(2):34-38.
13. Verma M, Rajput M, Sahoo SS, Kaur N. Neck circumference: Independent predictor for overweight and obesity in adult population. *Indian J Community Med.* 2017;42:209-213.
14. Kumar S, Gupta A, Jain S. Neck circumference as a predictor of obesity and overweight in rural central India. *Int. J. Med. Public Health.* 2012;2(1):62- 66.
15. Kelishadi R, Djalalinia S, Motlagh ME, Rahimi A, Bahreynian M, Arefirad T, et al. Association of neck circumference with general and abdominal obesity in children and adolescents: The weight disorders survey of the CASPIAN-IV study. *BMJ Open.* 2016;1-10.
16. Dayyat E, Kheirandish-Gozal L, Sans Capdevila O, Maarafeya MMA, Gozal D. Obstructive sleep apnea in children: Relative contributions of body mass index and adenotonsillar hypertrophy. *Chest* 2009;136(1):137-144.
17. Qaseem A, Dallas P, Owens DK, Starkey M, Holty JC, Shekelle P, et al. Diagnosis of obstructive sleep apnea in adults: A clinical practice guideline from the american college of physicians. *Ann Intern Med* 2014;161(3):210-220.
18. Corlateanu A, Covantev S, Botnaru V, Sircu V, Nenna R. To sleep, or not to sleep-that is the question, for polysomnography. *Breathe.* 2017;13:137-140.
19. Corlateanu A, Botnaru V, Sircu V, Covantev S, Montanari G. Obstructive sleep apnea and type 2 diabetes: Dual interaction. *Curr Respir Med Rev.* 2015;11: 292-298.

© 2018 Otoghile et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/46651>