



Liquidambar formosana Hance: A Mini-review of Chemical Constituents and Pharmacology

Xi Lin Ouyang^{1*}, Shan Yi¹, Hua Yuan Lu¹, Shang Mei Wu¹
and Hong Qiong Zhao¹

¹Youjiang Medical University for Nationalities, Chengxiang Road 98, Baise 533000, Guangxi, P.R. China.

Authors' contributions

This work was carried out in collaboration between all authors. Author XLO designed the study, managed the analyses of the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors SY and HYL searched the references and proof read the draft proposal and manuscript. Authors SMW and HQZ proof read the draft proposal and manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJMP/2016/29440

Editor(s):

(1) Marcello Iriti, Professor of Plant Biology and Pathology, Department of Agricultural and Environmental Sciences, Milan State University, Italy.

Reviewers:

- (1) Isiaka A. Ogunwande, Lagos State University, Ojo, Nigeria.
(2) Israel O. Obaroh, Kebbi State University of Science and Technology, Aliero, Nigeria.
(3) Anonymous, Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Bulgaria.
(4) Marisa Alves Nogueira Diaz, Federal University of Viçosa, Brazil.

Complete Peer review History: <http://www.sciencedomain.org/review-history/16704>

Mini-review Article

Received 10th September 2016
Accepted 22nd October 2016
Published 28th October 2016

ABSTRACT

Aims: *Liquidambar formosana* Hance is of great an ornamental and economic arbor. Its' various organs, such as the gum, fruits, barks, and leaves have a great medicinal value and commonly used in the pharmaceutical, food and cosmetic industries. Currently, several studies aimed at isolation and identification of active constituents of *L. formosana* and its pharmacological actions of different plant parts based on the treatment of several diseases. The main objective of this paper is to review recent advances of *L. formosana* in chemical constituents and pharmacology during the last two decades, in an attempt to provide useful references for plant drug studies.

Methodology: Information on *L. formosana* from journals and books published during the last two decades was collected based on worldwide accepted scientific databases via an electronic search (PubMed, Elsevier, Google Scholar, Springer, Web of Science and CNKI).

Results: The detailed phytochemical composition with respect to the pharmacological properties of

*Corresponding author: E-mail: oyxl2010@yahoo.com;

different parts of this multipurpose tree. Trace elements, volatile oils, terpenoids, flavonoids, phenolic acids and phenylpropanoids constituents are the natural plant secondary metabolites known from the different organs of *L. formosana*. The main pharmacological actions of *L. formosana* and compounds isolated therefrom include antitumor, antithrombotic, antimicrobial, antiviral, anti-inflammatory antidepressant, and antioxidant actions. *L. formosana* leaves are a strong antioxidant source and may either mitigate or prevent generation of free radicals.

Conclusion: Many pharmacological activities appear to be attributable to terpenoid and the flavonoid constituents; terpenoids had been proposed as being responsible for the antitumor, antimicrobial, and antiviral activities that had been documented for *L. formosana*. Based on the foundation of chemical constituents, their possible contribution to the demonstrated efficacy of extracts obtained from *L. formosana*, as well as new directions on quality control and application of this plant are suggested. These biologically active compounds are also being determined quantitatively by High Performance Liquid Chromatography (HPLC) analysis, which can be used to control the quantity of this plant. *L. formosana* will be further developed as medicinal and economic values, and thereby promote its cultivation. Further research is required to isolate and identify more new compounds contribute to the pharmacological effects.

Keywords: *Liquidambar formosana* Hance; chemical constituents; pharmacological properties; research progress.

1. INTRODUCTION

Plants of the genus *Liquidambar* as traditional Chinese and Ayurvedic medicines are dispersed mainly in Southeast Asia and America, but they have been cultivated in many countries around the world as ornamental plants. The genera contain four species, that is, *L. styraciflua* (known as American sweet gum), *L. orientalis* Mill. (Oriental sweet gum), *L. formosana* Hance (Chinese sweet gum) and *L. acalycina* H.T. Chang (Chang's sweet gum) [1]. *L. formosana* Hance (also known as maple), one of the genus *Liquidambar*, is a tall deciduous tree widely distributed in various regions of the south of the Qinling Mountains and Huaihe River in China, and also found in northern Vietnam, Laos and South Korea. *L. formosana* is a famous ornamental plant for leaves are green in spring and summer, and red in autumn. Different plant parts of *L. formosana*, such as the leaf, fruit, bark, and resin, are proved to be the treasures as natural medicinal plant resources (Fig. 1). Its leaves have the effects of relieving pain, detoxification, and hemostasis. It can be used for the treatment of acute gastroenteritis, dysentery, postpartum wind, infantile tetanus, carbuncles hair back, and so on. Its fruits (also known as Lulutong, Jiukongzi, *Fructus liquidambaris*), have been used as a traditional Chinese medicine (TCM) in China for thousands of years. It can treat joint arthralgia, numbness cramps, edema, puffiness, and so on. Aromatic essential oils from *L. formosana* resin (known as balsam) had been widely used for both medicinal and cosmetic purposes. This resin as an expensive TCM has

been used traditionally for the treatment of flutter injury, ulcer throat, vomiting, epistaxis, and trauma bleeding in China [2].



Fig. 1. *L. formosana* and its' various organs

As a kind of natural pigment resources, *L. formosana* has great potential in foods and cosmetics industry. Leaves of *L. formosana* have been taken as a rice dye for festival food by a small group of people in Guangxi province, a southern province of China. Moreover, *L. formosana* can be taken advantage of landscaping and xerogardening as an ornamental plant, and improve the soil structure in ecology [3]. Therefore, *L. formosana* has a good potential prospect for development because of its' medicinal, economic, ecological and ornamental values.

However, the shape of *L. formosana* resin is similar to that of pine rosin, which is often given the opportunity for unscrupulous traders to misdescribe two resins. In addition, as the big culture country of *L. formosana*, there are a lot of leaves, fruits, and barks as the offal abandon in the field but not to fully utilize every year in the rural area. So far no comprehensive review has been compiled from the literature encompassing the efficacy of *L. formosana*. Its versatile utility as a medicine, food additive, nutraceutical, and ornamental function motivated us to bridge the information gap and to form a comprehensive review on the chemical constituents identified from this plant and their divergent pharmacological properties.

2. CHEMICAL CONSTITUENTS

2.1 Trace Elements

L. formosana are rich in essential trace elements. The contents of magnesium (Mg), manganese (Mn), and calcium (Ca) in wild *L. formosana* leaves are very high compared to black rice, black beans and black sesame seeds, nearly 3 times higher for Mg, 13 times higher for Mn, and 115 to 400 times higher for Ca, but its contents of Zinc (Zn) and copper (Cu) are the same compared to these black nutritious food [4,5]. The leaf is the highest nutrient element in different organs, while the trunk had the lowest [6]. Therefore, *L. formosana* leaves can be used as medicinal plants and trace elements in food additives. Moreover, *L. formosana* have a strong enrichment capability for Cu, Zn, Mn, rare earth elements and other ions in soil [7], which are the material basis of blood glucose and blood pressure in medicinal values.

2.2 Volatile Oils

L. formosana contains a large number of volatile oils that can be obtained from the leaf and fruit of this plant by steam distillation. Its typical essential oil constituents, such as terpenes, cinnamic acids and small amounts of monoterpenes ingredients, including α -pinene, β -pinene, 4-terpineol, α -terpineol, camphene, juniperene, terpinene (including α , β , γ -configuration) are present [8,9]. Sesquiterpene ingredients including vitispirane, valeranal, valerenone are also found.

So far numerous studies have been conducted on different parts of *L. formosana* (Table 1).

Remarkable differences are found in the main components and the relative contents of the essential oils among the different studied organs, indicating that the pharmacological effects of the various parts have significant differences. Moreover, volatile oil component of *L. formosana* and *L. styraciflua* indicates a closer relationship based on GC/MS methods of qualitative and quantitative analysis, but the genetic relationship between *L. storax* [10]. Thus, leaves and resins from *L. formosana* were proved the potential sources for volatile oil. Differences of various organs, regional and species are the key factors in research and development of volatile oil.

2.3 Terpenoids

Terpenoids are the main chemical composition of *L. formosana*. A wide variety of compounds have been extracted and purified from the barks, resin of *L. formosana*, including diterpenes (such as iridoid and its derivatives) and triterpenoids (Table 2). The barks contain four iridoid glycosides (compounds **1-4**), which was often regarded as important active ingredients in medicinal plants [11]. A recent report showed that thirteen diterpenes (compounds **5-17**) were isolated from *L. formosana* resin [12]. Typical structures of triterpenoids, also contain a range of normal tetracyclic (dammarane, tirucallane) and pentacyclic (lupane, oleanane) triterpenoids are the most abundant components widely dispersed various organs of *L. formosana*. More than fifty triterpenes were isolated from different parts of *L. formosana* (Fig. 2).

2.4 Phenolic Acids and Tannins

Phenolic acids and tannins are one of the main components in leaves, branches, and barks of *L. formosana*. The composition of the phenylpropanoids of leaf mainly consists of gallic acid (**69**), p-hydroxy-benzoic acid (**70**), 3-methoxy-4-hydroxy-benzoic acid (**71**), 3,5-dihydroxy-4-methoxy-benzoic acid (**72**), 3,4-dihydroxy-benzoic acid (**73**), and 3,4-dihydroxy-5-methoxy-benzoic acid (**74**) [22]. Gallic acid has good biological activity widely distributed in plants and foods [27], it was often being used to evaluate the quality of this herb for its considerable contents in leaves and fruits of *L. formosana* [28]. It is reported that tannin content in leaf is nearly 13%, much higher than that of the bark [29], whereas *L. formosana* branch has the lowest concentration level of tannin [30].

2.5 Flavonoids

Flavonols and dihydro flavonols are the main existence forms of flavonoids in *L. formosana*. Several flavonoids, such as rutin (**75**), isoquercitrin (**76**), hyperin (**77**), trefoil bean glycosides (**78**), astragalin (**79**), catechins (**80**), epigallocatechin (**81**) were obtained from *L. formosana* (Fig. 3). In recently years, (2S)-5,7,4'-trihydroxyflavan-7-O- β -D-glucopyranoside (**82**), (2S)-5,7,4'-three hydroxy-5-O- β -D-glucopyranoside (**83**) and other new flavan glycosides were found in *L. formosana* leaf [31]. Flavonoids are also used as a class of common pigments with good antioxidant activity widely used in industry. The impact of various factors on total flavonoids extraction rate was determined for optimum extraction technology of total flavonoids [32]. Moreover, dynamic changes of flavonoids content were measured in different leaf during various collection periods. The optimal harvest time of *L. formosana* leaves was determined based on the contents of flavonoids

with different collection time [33]. Our study found that mature leaves as flavonoid-rich additives are the best resources of natural food pigments for the extraction in food technology [34].

2.6 Other Compounds

Three phenylpropanoids, that is cinnamic acid cinnamic ester [19], bornyl cinnamate, cinnamyl cinnamate [20], were isolated and identified from *L. formosana* resin. Furthermore, *L. formosana* leaf has been reported to contain a fairly unique group of constitutes called pigments, namely red pigment and melanin [35-37], which were commonly used as safe additives in the food industry. (7S,8S) -3-methoxyl-3'-O- β -D-glucopyranosyl-4':7,5':8-diepoxyneolignan-4,9'-diol [31], choline, acetylcholine, glucuronidase, β -sitosterol, β -sitosterol glucoside, 5-formylbilinone have also been isolated from the leaf of *L. formosana*.




Fig. 2. Structures of selected diterpenes and triterpenoids from *L. formosana*

Table 1. Analysis of volatile oils from different parts of *L. formosana*

Organs	Methods	Constituents	References
1 Leaf	GC/MS/DS	Twenty compounds were identified, accounting for 84.88%. 4-terpineol (27.17%), Δ 1(2), 8-menthadiene (9.39%), β -caryophyllene (7.06%) and 7-hydroxycoumarin (6.82%) are the main ingredient.	[13]
2 Leaf	GC/MS	Forty-seven compounds were identified, accounting for 98.11% of naphtha, including 30 terpenoids, 14 aliphatic compounds and 3 aromatic compounds. The main components are β -pinene (21.18%), α -pinene (20.70%), and (E)-2-hexenal (7.64%), limonene (7.59%), β -caryophyllene (6.08%), etc.	[14]
3 Leaf	GC/MS	Twenty-nine compounds were identified, accounting for 81.04% of the total essential oil. The main components are <i>n</i> -palmitic acid (27.03%) and 9,12,15-shap acid (13.35%).	[15]
4 Fruit	GC/MS	Twenty-nine compounds were identified. The main components compose of β -pinene (16.69%), 1S- α -pinene (15.22%) and caryophyllene (12.54%).	[16]
5 Leaf	GC/MS	25 compounds were identified, accounting for 92.81% of total volatile oil, which contains 18 terpenes, 5 fatty substance and 2 aromatic substances. The main components were composed of α -pinene (34.48%), β -pinene (19.25%), and limonene (26.97%).	[17]
6 Leaf, bark, heartwood	GC/MS	Fifteen components were identified from leaf, seven components from the bark, and nine components from heartwood.	[18]
7 Resin	GC/MS	Volatile oil in resin is 12%. The main components are α -pinene (24.92%), β -pinene (23.62%), camphene (8.79%), terpinolene (6.41%), bornyl acetate (2.97%) and β -caryophyllene (19.62%). The content of caryophyllene is nearly 20%.	[19]
8 Resin	GC/MS	α -pinene (23.3%), β -caryophyllene (22.7%) and β -pinene (19.6%) are the main ingredient.	[20]
9 Leaf and stem	GLC/FID and GLC/MS	Sixty-four components were identified from the leaves and stems. The major components of the leaf were d-limonene (22.34%), α -pinene (27.95%) and β -pinene (11.20%), and that of the stem oil were germacrene D (10.91%), α -cadinol, d-limonene (12.89%), α -pinene (14.19%), and β -pinene (5.34%).	[1]

Table 2. The reported terpenoids isolated from *L. formosana*

No.	Names	Sources	References
1	Monotropein	Barks	[11]
2	Monotropein methyl ester		
3	6 α -hydroxy geniposidic		
4	6 β -hydroxy geniposidic		
5	Liquidambolide A	Resin or	[12]
6	Liquiditerpenoic acid A	balsam	
7	Liquiditerpenoic acid B		
8	15-hydroxydehydroabietic acid		
9	15-hydroxy-7-oxodehydroabietic acid		
10	2 α -hydroxydehydroabietic acid (10)		
11	12-hydroxy-7-oxo-5,8,11,13-tetraene-18,6-abietanolide		
12	Picealactone A		
13	12 α -hydroxyabietic acid		
14	12 β -hydroxyabietic acid		
15	15-hydroxy-6-enedehydroabietic acid		
16	Aquilarabietic acid K		
17	Pimaric acid		
18	2 α ,3 α -dihydroxyolean-12-en-28-al		[20]
19	3 α -hydroxyolean-12-en-30-ol		
20	3 α -hydroxyolean-2-oxo-12-en-28-al		
21	3 α ,28-dihydroxyolean-12-ene		
22	28-hydroxy- β -amyrone		
23	Oleanonic aldehyde		
24	3-oxoolean-12-en-28-oic acid		
25	3-oxoolean-12-en-28-yl acetate		
26	3 α ,25-dihydroxyolean-12-en-28-oic acid		
27	3,28-dioxobetulin		
28	3-oxobetulin		
29	Betulonic acid		
30	3-oxoursa-12-en-28-al		
31	Liquidambronic acid		[21]
32	Hmbronic acid		
33	Ambronic acid		
34	Formosolic acid		
35	Ambradiolic acid		
36	Liquidambronal		
37	Ambronal		
38	Oleanolic ketone alcohol		[19]
39	3-oxo-olean-12-ene-28-oic acid (oleanonic acid)		
40	3-oxo-lup-20(29)-ene-28-oic acid (liquidambronic acid)		
41	3 β ,23,29-trihydroxy-olean-12-en-28-oic acid- β -D-glucopyranosyl ester	Leaves	[22]
42	3 β ,6 β -dihydroxylup-20(29)-en-28-oic acid β -glucopyranosyl ester	Roots	[23]
43	2 α -acetoxyl-3 β ,6 β -dihydroxylup-20(29)-en-28-oic acid β -glucopyranosyl ester		
44	Stigmast-4-en-3-one		
45	(24R)-3 β -hydroxy-24-ethylcholest-5-en-one		
46	β -amyrin		
39	Oleanonic acid		
47	Arjunic acid		
48	Hederagenin 28-O- β -D-glucopyranoside		
49	(2 α ,3 β ,4 α)-23-(acetyloxy)-2,3-dihydroxy-olean-12-en-28-oic acid β -D-glucopyranosyl ester		
50	Arjunglucoside II		

No.	Names	Sources	References
51	Quadranoside I		
52	2,4,6-trimethoxyphenol-1-O-β-D-glucopyranoside		
53	3,30-di-O-methyllellagic acid-40-O-β-dxylopyranoside		
54	3,4,5-trimethoxyphenyl-(6-O-galloyl)-O-β-D-glucopyranoside		
55	3,4,5-trimethoxyphenyl-6-O-syringoyl-β-D-glucopyranoside		
56	3-oxo-11α,12α-epoxyoleanan-28, 13β-olide	Fruits	[24]
57	3-oxo-12α-hydroxy-oleanan-28, 13β-olide		
58	3α-acetoxyl-25-hydroxyoleanan-12-en-28-oic-acid		
39	Oleanonic acid		
59	Ursolic acid		
60	2α,3β,23-trihydroxyolean-12(13)-en-28-oic acid (arjunolic acid)		[25]
61	2α,3β-dihydroxy-23-norolean-4(24),12(13)-dien-28-oic acid		
62	11α-methoxyl-28-nor-β-amyrenone		[26]
63	28-nor-β-amyrenone		
64	3-oxo-12α-hydroxy-oleanan-28, 13β-olide		
65	3α-acetoxyl-25-hydroxy-olean-12-en-28-oic acid		
66	Erythodiol		
67	Betulonic acid		
68	Styracin		




Fig. 3. Structures of selected flavonoids from *L. formosana*

3. PHARMACOLOGICAL ACTIVITIES

3.1 Antitumor and Anticancer

Petroleum ether and dichloromethane extracts of Lulutong exhibited significantly cytotoxic on human colon adenocarcinoma cells HT-29, whereas methanol extract showed a minimal to negligible effect on cytotoxicity [38]. It has been found that 25-acetyl-3α-hydroxy oleanolic 12-en-28-oic acid and 3α, 25-dihydroxy-12-ene oleanolic acid, two triterpenic acids isolated from Lulutong showed significant inhibitory effects on HT-29 and HCT-116 cells. Another two oleanane triterpenoid acids, 3α-acetoxy-25-hydroxy-olean-12-en-28-oic acid and lantanolic acid isolated from the MeOH extract of the fruit by activity-guided fractionation, had been reported as NFAT transcription factor regulators. These compounds were proven to be a potent antitumor promoting activity using an in vitro assay on NFAT with IC₅₀ of 4.63 μM (3α-acetoxy-25-hydroxy-olean-12-en-28-oic acid) and 12.62 μM (lantanolic acid), respectively [39]. On the other hand, among

these pentacyclic triterpenes, olean triterpenes with 3-keto skeleton showed significant cytotoxicity against MDA-MB-435S tumor cells [40], suggesting that the 3-keto group is a critical structural factor for activity. The result indicated that triterpenic acids are a class of important antitumor components widely distributed in this plant.

3.2 Antithrombotic Activity

The extracts of *L. formosana* resin and volatile oil at the dose of 1.0 mg/mL can obviously inhibit thrombus formation in all treated animals. Its antithrombotic effect was related to promoting plasmin activity and increase the level of cAMP on platelet, indicating the extracts of *L. formosana* resin had an antithrombotic effect [41]. The recalcification experiment shows that volatile oil from leaves has also a good coagulation effect *in vitro* [42]. 28-oleanolic acid type carboxyl triterpenoids compounds from the extracts of maple and petroleum ether significantly inhibited ADP-induced platelet aggregation, suggesting that pentacyclic

triterpenoids have anti-platelet aggregation activity [40].

3.3 Antimicrobial and Antiviral Activities

The extracts of *L. formosana* leaves are commonly used as an antibacterial material in traditional folk medicine in China. *L. formosana* leaves showed strong effects against *Staphylococcus aureus*, *Staphylococcus*, *Shigella flexneri*, typhoid common pathogens and opportunistic pathogens *in vitro* antibacterial activity. However, little inhibition effects were found on *Escherichia coli* and *Candida albicans* [43]. Essential oil from *L. formosana* leaves have also a good stability in the acidic environment, which demonstrated strong inhibition of gram-positive bacteria and fungi, such as *Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli*, *Aspergillus flavus* and *Penicillium*, but relatively weak against Gram-negative bacteria [44]. Several compounds, such as 3 α ,25-dihydroxy-12-en-28-oic acid and bornyl cinnamic acid from the essential oils of *L. formosana*, were found to be responsible for the antibacterial effects on *Lenzites betulina* and *Laetiporus sulphureus* [20].

A recent report showed that the extract of *L. formosana* may be applicable as an inhibitor of Neuraminidase (NA, as sialidase influenza virus) on influenza virus A/California/7/2009 (H1N1) NYMCNA test *in vitro* [45]. It can reduce the replication of the virus, and reduce cytopathic effect (CPE) and the amount of virus induced by the MDCK cells when the concentration of the extract was at the dose of 25-250 $\mu\text{g/mL}$.

3.4 Anti-inflammatory Activity

Activated NFAT can promote the migration and invasion of cancer cells. However, excessive activated NFAT will cause a series of pathological and inflammatory reactions, such as autoimmunity and immune rejection. Dat et al [39] found that oleanane-type triterpenoids from the methanol extract of Lulutong have a strong inhibitory activity of NFAT. An aqueous decoction of Lulutong has anti-inflammatory and analgesic effects to the rat's paw edema induced by yeast. Furthermore, the essential oil obtained from *L. formosana* leaves was also demonstrated to exhibit anti-inflammatory activity in BEAS-2B human lung by detecting the expression of IL-6 protein used dexamethasone + TNF- α -induced cells as a positive control group (Dex) in epithelial cell inflammation model [42]. The underlying mechanisms for anti-inflammatory

activity include reducing nitrite oxide generation, secretion levels of TNF- α and IL-6, and expression levels of prointerleukin- β , inducible nitric oxide synthase, and cyclooxygenase-2 in lipopolysaccharide (LPS)-activated mouse macrophages [9].

3.5 Antidepressant Activity

Monoamine oxidase (MAO) is a flavin adenine dinucleotide (FAD), containing enzyme of the outer mitochondrial membrane. The organic acid extracts of *L. formosana* showed a strong inhibition activity to MAO with inhibition at $\text{IC}_{50} < 10 \mu\text{g/mL}$, suggesting good antidepressant activity of *L. formosana* [46]. It was also found that the total phenolic extracts of the fruits of *L. formosana* strongly inhibited MAO with an IC_{50} value of 5.9 $\mu\text{g/mL}$. Further bioactivity-guided chromatographic fractionation led to the isolation of gallic acid, ethyl gallate, p-digalloyl acid, and ethyl-p-digallate. These compounds showed high inhibitory activities *in vitro* against rat liver MAO with IC_{50} values of 1.7, 1.9, 1.5, and 1.1 $\mu\text{g/mL}$, respectively, whereas the standard IC_{50} value of iproniazid was 1.8 $\mu\text{g/mL}$ [47]. So, the fruits of *L. formosana* can be used as anti-depressant drugs levity.

3.6 Antioxidant Activity

L. formosana has been reported to exhibit other diverse activities, such as antioxidant activity. The extracts of *L. formosana* leaf showed good antioxidant activities than butylated hydroxytoluene (BHT), a synthetic antioxidant. The order of leaf extract with oxidation resistance is ethanol extracts > aqueous extract > acetone extract [48]. A methanol extract of *L. formosana* fruits also exhibited strong antioxidant activity, whereas petroleum ether and dichloromethane fractions had minimal or negligible effects [38].

Furthermore, the extract of *L. formosana* leaf can significantly improve the phagocytic percentage and phagocytic index of peritoneal macrophages in mice, suggesting a good enhanced non-specific immunity and cellular immunity effect [49].

4. FUTURE PROSPECTS

L. formosana is regarded as an important medicinal and ornamental plant with various economic, medicinal and ornamental benefits and ecological potential. Although preliminary studies of *L. formosana* are under way in

different laboratories to use the antitumor, antithrombotic, anti-inflammatory, antimicrobial, antiviral and antidepressant activities, these studies should be extended to humans in view of the medicinal nature of the plant. Moreover, diverse pharmacological activities of *L. formosana* are attributed to phytochemical compositions, especially volatile oils, phenylpropanoids, flavonoids and terpenoids rich in leaves, resins, and fruits, which are responsible for the effects of this medicinal plant. Most of the terpenoids identified in this paper were obtained from the extracts of resins, roots, and fruits. Most of the flavonoids were found from the leaves. Phenylpropanoids exist widely in various parts of *L. formosana*. Terpenoids, flavonoids, and phenylpropanoids as three kinds of main active constituents are also widely used as reference compounds to determine medicinal plants quality.

Food additives obtained from natural product have become the most important way for humans to enrich food resource. *L. formosana*, known to contain trace elements, pigments, phenylpropanoids, and flavonoids with noteworthy antioxidant activity, may be a good candidate as food additives. However, the volatile oils composition of *L. formosana* is strongly influenced by both different parts and geographical origins factors, which were determined by GC-MS.

As a folk medicine, different parts of *L. formosana* have its medicinal values. Resin and fruits as two important TCMs recorded in "Chinese Pharmacopoeia" attract more attentions than other organs. *L. formosana* resin is rich in essential oils and has been developed as a spice. It is noted that resin as a relatively scarce and expensive resource, should have further deeply research in terms of quality control and evaluation, such as identification between *L. formosana* resin and pine rosin.

In summary, although numerous studies have been conducted on different parts of *L. formosana*, but there is a dire need to be carried out in isolation and identification of new compounds from different parts of this plant in the near future. With a suitable pharmacological model, more studies will gradually screen the main active ingredients or the pure compounds from the crude extracts of *L. formosana*. Furthermore, these biologically active compounds can be determined quantitatively by HPLC analysis, which can be used to control the

quantity of this plant. *L. formosana* will be further developed as medicinal and economic values, and thereby promote its cultivation.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

The authors thank the Scientific Research Fund of Guangxi Provincial Education Department (No. YB2014295) and the open fund of the Key Laboratory of Youjiang Basin Authentic Traditional Chinese Medicine (National Herbs) of Guangxi Universities (No. szdsysyy2015301) for financial support.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. El-Readi MZ, Eid HH, Ashour ML, Eid SY, Labib RM, Sporer F, et al. Variations of the chemical composition and bioactivity of essential oils from leaves and stems of *Liquidambar styraciflua* (Altingiaceae). J Pharm Pharmacol. 2013;65(11):1653-1663.
2. China Pharmacopoeia Commission. Pharmacopoeia of the People's Republic of China. Beijing: People's Medical Publishing House. 2015;356-357.
3. Wang G, Tian D, Yan W, Zhu F, Li S. Impact of litter addition and exclusion on soil respiration in a *Liquidambar formosana* forest and a nearby *Cinnamomum camphora* forest of central southern China. Acta Ecologica Sinica. 2009;29(3):1607-1615.
4. Huang L, Shong XP. Studies on the chemical constituents of hainan island *Liquidambar formosana*. Stud Trace Elem Health. 2001;18(1):45.
5. Zhong YT, Wang L, Wang XL, Sun XT. Determination of Gannan wild maple leaves trace elements by flame atomic absorption spectrometry. Lishizhen Med Mater Medica Res. 2010;21(6):1457-1458.

6. Yang LL, Wen SZ, He GX. Biological cycling of nutrients in *Liquidambar formosana* plantation in suburban are of Changsha. J Fujian Coll For. 2012;32(1): 48-53.
7. Huang L, Song XP. Studies on the medicinal function and medicinal taste on the *Liquidambar formosana*. Food Sci. 2002;23(5):132-134.
8. Lee Y-S, Kim J, Lee S-G, Oh E, Shin S-C, Park I-K. Effects of plant essential oils and components from Oriental sweetgum (*Liquidambar orientalis*) on growth and morphogenesis of three phytopathogenic fungi. Pestic Biochem Physiol. 2009;93(3): 138-143.
9. Hua KF, Yang TJ, Chiu HW, Ho CL. Essential oil from leaves of *Liquidambar formosana* ameliorates inflammatory response in lipopolysaccharide-activated mouse macrophages. Nat Prod Commun. 2014;9(6):869-872.
10. Chen YD, Hu ZD, Gu Y. Study on chemical composition on flavonoids and terpenes in *Liquidambar*. Chem Ind For Prod. 1991;11(2):157-164.
11. Jiang ZH, Zhou RH, Kouno I. Iridoid components from *Liquidambar formosana* bark. Chin Tradit Herbal Drugs. 1995; 26(8):443-444.
12. Shang HJ, Li DY, Wang WJ, Li ZL, Hua HM. Three new diterpenoids from the resin of *Liquidambar formosana*. Nat Prod Res. 2014;28(1):1-6.
13. Jiang ZH, Zhou RH. Chemical constituents of volatile oil from leaves of *Liquidambar*. Chin Herbal Med. 1991;14(8):34-35.
14. Liu YM, Liu YM, Li PX, Guo Y, Ma M. Extraction process and component analysis of volatile oil from *Liquidambar formosana* leaves. Chem Ind For Prod. 2009;29(4):77-81.
15. Chen HS, Zhao ZG, Liang XY, Jiang QY, Jiang XH. Studies on chemical constituents of volatile oil from *Liquidambar acalycina*. Guihaia. 2009; 29(1):136-138.
16. Cai AH, Zhao ZG, Chen HS, Jiang XH, Yan XQ. Analysis of volatile constituents from the fruit of *Liquidambar formosana* and *L. acalycina* by GC-MS. J Guilin Univ Technol. 2012;35(2):245-249.
17. Zhang YH, Feng J, Jin XH, Yao XQ, Yang ZJ. Components analysis of volatile oil from *Liquidambar formosana* leaves by GC-MS. Chin J Exp Tradit Med Form. 2012;18(22):81-83.
18. Shang QW, Ma L. Research on volatile oils North America *Liquidambar formosana*. Nei Mongol J Traditional Chin Med. 2013; 32(26):30-30.
19. Liu H, Chen MY. Studies on the chemical constituents of *Liquidambar formosana* resin. Chem Ind For Prod. 1995;15(3): 61-66.
20. Chien SC, Xiao JH, Tseng YH, Kuo YH, Wang SY. Composition and antifungal activity of balsam from *Liquidambar formosana* Hance. Holzforschung. 2013; 67(3):345-351.
21. Liu C, Xu JF, He QM, Yu QS. Chemical components of the *Liquidambar formosana* H. Resin. Chin J Org Chem. 1991;11(5): 508-510.
22. Liao YY, Fu HZ, Zhou GP, Zhong RJ, Yuan H, Guo Q, et al. Chemical constituents from leaves of *Liquidambar formosana*. J Chin Med Mater. 2014; 37(12):2219-2221.
23. Yu J, Liu SS, Xuan LJ. Two new lupene-type triterpenoids from the roots of *Liquidambar formosana*. Nat Prod Res. 2012;26(7):630-636.
24. Li C, Sun YR, Sun YF. Chemical composition of *Fructus liquidambaris*-Lulutong. Acta Pharm Sinica. 2002;37(4): 263-266.
25. Lai ZQ, Dong Y. Studies on the chemical constituents of the lulutong(l)-structure determination of a novel pentacyclic triterpene. Acta Sci Nat Univ Sunyatseeni. 1996;35(4):64-69.
26. Shang HJ, Wang WJ, Li DY, Hua HM, Li ZL. A new triterpenoid isolated from fruits of *Liquidambaris fructus*. Chin Tradit Herbal Drugs. 2014;45(9):1207-1210.
27. Cirillo G, Kraemer K, Fuessel S, Puoci F, Curcio M, Spizzirri UG, et al. Biological Activity of a Gallic Acid-Gelatin Conjugate. Biomacromolecules. 2010;11(12):3309-3315.
28. Feng Y, Zhang YY, Wu SM, Lu YM, Lan WQ, Ouyang XL. Extraction of gallic acid in *Fructus liquidambaris* from Guangxi by orthogonal experiment. Central South Pharm. 2015;13(1):21-23.
29. Zheng Y, Zhang Q. Extraction and complexometric titration analysis of tannin in the barks and leaves of *Liquidambar formosana* Hance. Jiangxi Chem Ind. 2004;2:99-102.
30. Zhang Q, Zheng Y. The organic solvent extracts tannins in the branches and leaves of *Liquidambar formosana* Hance

- and the content is determined. Jiangxi Chem Ind. 2006;3:71-74.
31. Zhong RJ, Yuan H, Fu HZ, Zhou GP, Wu X, Zhang CH, et al. Three new compounds from the leaves of *Liquidambar formosana*. J Asian Nat Prod Res. 2013;15(12):1249-1255.
 32. Li Y, Song GZ. Ultrasonic-assisted extraction of total flavonoids from *Liquidambar formosana*. J Shanxi Agric Sci. 2010;38(8):69-70.
 33. Liu YM, Liu YM, Ma M, Li CX, Llan CX, Cao SJ. Extraction and dynamic content of total flavonoids in leaves of *Liquidambar formosana* Hance. Food Sci. 2010;31(4):35-38.
 34. Lu HY, Wu SM, Yi S, Ouyang XL. Comparison of total flavonoid content leaves of *Liquidambar formosana* at different age. Hubei Agric Sci. 2016; 55(16):4203-4205, 4213.
 35. Fang YM, Zhang P, Wang YH, Tan P, Wang P. Study on extraction of melanin from leaves of guizhou *Liquidambar formosana* hance. Hubei Agric Sci. 2015; 54(13):3230-3233.
 36. Jiang XL. Extraction of red pigment from *Liquidambar formosana* leaves. China Brewing. 2011;30(2):139-142.
 37. Sheng XY, Jun ZX, Rong ZL. The analysis of some water-soluble microelements in maple leaves. J Guangxi Teachers Coll. 2004;17(3):5-7.
 38. Zhang J, Chou G, Liu Z, Koh GY. In vitro cytotoxicity and antioxidation of a whole fruit extract of *Liquidambar formosana* exerted by different constituents. Eur J Med Plants. 2015;6(1):34.
 39. Dat NT, Lee IS, Cai XF, Shen G, Kim YH. Oleanane triterpenoids with inhibitory activity against NFAT transcription factor from *Liquidambar formosana*. Biol Pharm Bull Mar. 2004;27(3):426-428.
 40. Yang NY, Chen JH, Zhou GS, Tang YP, Duan JA, Tian LJ, et al. Pentacyclic triterpenes from the resin of *Liquidambar formosana*. Fitoterapia. 2011;82(6):927-931.
 41. Zhu L, Guo JX. Antithrombotic effects of *Liquidambar formosana* resin and volatile oil. Chin Tradit Herbal Drugs. 1991;22(9):404-405.
 42. Yang ZJ, Zhang YH, Feng J, Yao XQ, Jin XH. Pharmacological activity and chemical composition of *Liquidambar formosana* leaves volatile oil. Sci Technol Food Ind. 2013;34(4):76-79.
 43. Zhong YT, Wang XL, Ma LL. Studies on antimicrobial activity of *Liquidambar formosana* Hance leaf. Lishizhen Med Mater Medica Res. 2007;18(7):1693-1694.
 44. Liu YM, Liu YM, Li PX. Study on antimicrobial activities of essential oil from leaves of *Liquidambar formosana* Hance as well as its antioxidant activity. Food Sci. 2009;30(11):134-137.
 45. Tian L, Wang ZY, Wu H, Wang S, Wang Y, Wang YY, et al. Evaluation of the anti-neuraminidase activity of the traditional Chinese medicines and determination of the anti-influenza A virus effects of the neuraminidase inhibitory TCMs *in vitro* and *in vivo*. J Ethnopharmacol. 2011;137(1):534-542.
 46. Yang ZD, Ren J, Shu ZM. Monoamine oxidase inhibitory activity of the total alkaloid and organic acid from Chinese herbal medicines. Adv Mater Res. 2013; 781-784:899-902.
 47. Yang ZD, Liang JB, Xue WW, Sheng J, Shi Y, Yao XJ, et al. Phenolic compounds from *Liquidambar formosana* fruits as monoamine oxidase inhibitors. Chem Nat Compd. 2014;50(6):1118-1119.
 48. Wang K, Pan Y, Wang H, Zhang Y, Lei Q, Zhu Z, et al. Antioxidant activities of *Liquidambar formosana* Hance leaf extracts. Med Chem Res. 2010;19(2):166-176.
 49. Zhong YT, Wang XL, Sun XT, Xu J, Wang L, Zhang WP, et al. The study of extract from leaves of *Liquidambar formosana* Hance on immune-regulatory effects in mice. Pharmacol Clinics Chin Mater Med. 2012;28(1):124-126.

© 2016 Ouyang et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
 The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/16704>