



Role of Ultrasonography and Conventional Radiography in the Detection of Maxillofacial Fractures: A Comparative Study

Ayesha Kamran^{a#} and Wisam A. Razzak Al-Gorjia^{a*}

^a *D.H.Q. Sargodha (Affiliated with Sargodha Medical University), Pakistan.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i60A34546

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/78679>

Original Research Article

Received 15 October 2021
Accepted 19 December 2021
Published 20 December 2021

ABSTRACT

Objective: With the collaboration of the trauma department, our study was designed to compare the effectiveness of ultrasonography (USG) and conventional radiography in the detection of bony fractures related to oral and maxillofacial regions.

Methodology: This comparative study was conducted from March 2020 to March 2021 by the Radiology department of Sarghoda medical college hospital with the collaboration of the trauma department. Ultrasonography was performed by using GE- USG machine along with a linear extraoral transducer (frequency range 7-15 MHZ). Patients were asked to sit in a seated position facing the sinologist. Transducers were placed over the site by applying the sterile gel.

Results: The overall sensitivity and specificity rate of ultrasonography was reported as 83.33% and 98.88% respectively in all sites whereas the sensitivity and specificity rate of conventional radiographs were reported as 70.24%, 100%. The negative predictive value of USG was reported as 96.17% along with 94.59% positive predictive value. In the contrast, conventional radiography gave a better positive predictive value (100%) than USG In our study we found better results of ultrasonography in terms of sensitivity and negative predictive value.

Conclusion: In conclusion, our study depicts that ultrasonography is an economical, useful diagnostic tool for examining the bony fractures of facial trauma with a better sensitivity rate when compared to conventional radiographs.

[#]Associate Prof of Radiology,

^{*}Corresponding author: E-mail: wisamrazzak@gmail.com;

Keywords: Conventional radiographs; bony fracture; ultrasonography.

1. INTRODUCTION

Clinical diagnoses of facial injury are insufficient to examine the fracture lines which demand radiological assessment [1]. Conventional assessment of facial trauma involves a structural approach which includes history, palpation inspection, and auscultation [2]. Conventional radiographs play a key role in examining the location, magnitude, and displacement of fractures [3]. Nowadays the traditional method of conventional radiography is replaced with high imaging modalities including computed tomography and cone-beam CT (CBCT). However, these methods have some drawbacks in terms of expensive equipment, difficulty in maintaining a steady position, radiation exposure, and lack of availability in many regions of the world [1]. Hence, many physicians limit these methods and use ultrasonography in facial bony fractures. Ultrasonography is one of the best methods for the identification of pathology related to soft tissues present in the head and neck region [1]. In past, this method was widely used for detecting soft tissue lesions, salivary glands [2]. Very few studies reported its role in injuries related to maxillofacial [3]. Previous studies reported 85% accuracy of ultrasonography in detecting fractures related to zygomatic-orbital complex (ZMC) [1,4]. On contrary, another study reported the usefulness of ultrasonography in the visualization of the zygomatic arch and the frontal sinus anterior wall [5]. Ultrasonography failed to penetrate in deeper bony structure so its use is only restricted to examine the superficial facial fractures.

In many regions of midfacial fractures like orbit [6,7], nasal bone [8,9] and zygotic arch [10-12] ultrasonography reported successive outcomes. It also assists during surgery in examining the fracture reductions to attain adequate repositioning of fractured segments [13,14]. A systematic review conducted in 2011 reported the usefulness of ultrasonography in fractures related to the anterior maxillary wall in pregnant women and children without any radiation exposures [15]. With the collaboration of the trauma department, our study was designed to compare the effectiveness of ultrasonography and conventional radiography in the detection of bony fractures related to oral and maxillofacial regions.

2. METHODOLOGY

This comparative study was conducted from March 2020 to March 2021 at the Radiology Department of Sarghoda medical college hospital with the collaboration of trauma department. The sample size of 277 sites was estimated by predicting 0.05% error with expected 90% sensitivity. However, in the final analysis, 20 patients were recruited for screening. All the suspected patients of a facial skeleton who underwent CT scans and conventional radiographic examinations, visiting our facial and trauma center from March 2020 to March 2021 were included. To access the effectiveness of both techniques we set time frame for inclusion because in delayed diagnosis many symptoms related to soft tissue resolved. Patients who were diagnosed during 0-15 days of trauma were only included.

Only conscious and cooperative patients during screening were included. Patients suffering from severe soft tissue lacerations, oedema, and having complex fractures were excluded from the research. Patients with dressing and abrasions were also not included due to the probability of intense pain and discomfort. The study was carried out by following complete protocol for radiological assessment. Single examiner collected all the demographic details of patients including age, sex, occupation, trauma details including date and time of trauma, cause of injury, and treatment after trauma. The same examiner conducted a clinical examination of each patient and recorded relevant findings of extraoral and intraoral examinations. Standard techniques of conventional radiographs were used for submentovertex view, water's view, and a panoramic view. Interpretation of the radiographs was done by two senior radiologists. On the other hand, a CT scan was performed by using 64 slice CT scanner and their interpretation was done by a single radiologist. We considered the results of CT as a gold standard for comparison with other techniques. Ultrasonography was performed by using GE- USG machine along with a linear extraoral transducer (frequency range 7-15 MHZ). Patients were asked to sit in a seated position facing the sinologist. Transducers were placed over the site by applying the sterile gel. For this study, we define fracture as an interruption that occurs in the radiopaque line of the bony contour including displacement. The research was conducted by

following all the principles of Helsinki. Written consents were taken from patients and they were well aware of the objectives and consequences of research [16].

Statistical analysis was performed using SPSS 23.0. Frequencies were noted for all continuous variables. Sensitivity, specificity, negative and positive predictive values of USG and conventional radiographs were observed for comparison.

3. RESULTS

This study enrolled a total of 20 patients. 18 of the 20 patients were male, with an average age of 34.4 years (19-75 years). In 90% of cases, traffic accidents were the leading cause of injury, while 10% were caused by falls from height. Approximately, 15-20 minutes per patient were consumed for examining all fracture sites using ultrasonography. During USG examination not a single patient reported discomfort or pain. For the conventional radiographs, approximately 20 minutes were taken for each radiograph. CT scan was performed in an average timeframe of 30-40 minutes for each patient. Each patient was

bilaterally examined for 11 sites of the face. According to the standard protocol of CT scan, 84 fractured sites were examined. Out of these 84, fifty-nine sites were accurately detected by conventional radiographs whereas ultrasonography detected 74 sites. Out of these 74 sites four false-positive results were observed. Ultrasonography detected all the sites (mentioned in Table 1) accurately whereas conventional radiographs gave more accurate results for mandibular condyle/subcondyle sites. However, both methods failed to detect orbital floor fracture.

The overall sensitivity and specificity rate of ultrasonography was reported as 83.33% and 98.88% respectively in all sites whereas the sensitivity and specificity rate of conventional radiographs were reported as 70.24%, 100%. The negative predictive value of USG was reported as 96.17% along with 94.59% positive predictive value. In the contrast, conventional radiography gave a better positive predictive value (100%) than USG. In our study we found better results of ultrasonography in terms of sensitivity and negative predictive value (Table 2).

Table 1. Fracture distribution according to injury site [16]

Examination Sites	USG (True Positive + False Positive)	Sensitivity of USG	Specificity of USG	Radiographs (True Positive + False Positive)	CT Scan Gold standard
Condyle/Subcondyle	4 (4+0)	80%	100%	5 (5+0)	5
Zygomatic Arch	12 (12+0)	100%	100%	12 (12+0)	12
Angle	2 (1+1)	100%	97.43%	1 (1+0)	1
Fronto-Zygomatic Process	3 (3+0)	33.33%	100%	0 (0+0)	9
Symphysis/Parasymphysis	7 (7+0)	100%	100%	7 (7+0)	7
Zygomatic Bone	2 (2+0)	50%	100%	1 (1+0)	4
Orbital Floor	0 (0+0)	0%	100%	0 (0+0)	3
Anterior Wall of Frontal Sinus	4 (4+0)	100%	100%	0 (0+0)	4
Supraorbital Margin	5 (2+3)	100%	92.10%	2 (2+0)	2
Infraorbital Margin	19 (19+0)	100%	100%	17 (17+0)	19
Anterior/Lateral Wall of Maxillary Sinus	16 (16+0)	88.89%	100%	14 (14+0)	18

Table 2. Comparison of sensitivity and specificity rates of conventional radiographs versus ultrasonography [16]

Method		Gold Standard (CT scan)		Sensitivity	Specificity
		No fractures	Fractures		
Conventional radiographs	No fractures	356 (TN)	25 (FN)	70.24%	100%
	Fractures	0 (FP)	59 (TP)		
	No fractures	352 (TN)	14 (FN)		
USG	Fractures	4 (FP)	70 (TP)	83.33%	98.8%

4. DISCUSSION

Face fractures occur at any site so the early detection of the site is effective time management. These fractures may occur alone or with a combination of several other injuries which may cause severe complications at a later stage. CT scan has some drawbacks in the form of high radiation and high probability of developing cataracts limit used in children and pregnant women [3]. Furthermore, CT scanning is a too expensive and time taking procedure in many parts of the world when used in isolated simple fractures [3]. These drawbacks laid the foundation for developing new and better imaging techniques which at least reduced these risks. On contrary, ultrasonography has many advantages in terms of cheap method, without radiation exposure, fast and high availability but demands experienced physicians to handle or interpret [5]. Ultrasonography can also reveal the various phases of fracture including healing. With these advantages, USG has privileged the other conventional radiography. However, ultrasonography cannot penetrate deep bony structures [5]. With this drawback, its use is currently limited to evaluating the superficial structures. Although, resolution can be increased still the deep penetration cannot be acquired. In present study, total of 84 fractures sites out of 440 were observed according to the gold standard of CT scan. In our study ultrasonography showed 83.33% sensitivity and 98.88% specificity of all fractured sites when compared to the CT scan. In the present study 94.59% positive and 96.17% negative predictive value of ultrasonography were observed. When we compared these findings to those of other studies, it was discovered that none of them included all fractured sites. However, the sensitivity and specificity rates were similar to those studies which used ultrasonography for analyzing facial fractures. A study by Ayoub et al. reported an 85% accuracy rate of ultrasonography but their results vary at different sites in specificity, and positive predictive values

[1]. When examining each site separately we found 100% sensitivity and specificity of ultrasonography for the anterior wall of the frontal sinus, parasymphysis, and zygomatic arc when compared to the conventional radiography. These results depict that no false positive or false negative values were present at these sites. Our results also depicted that ultrasonography is more reliable at the lateral wall of the maxillary sinus with 94% sensitivity and 100% specificity rate. At the anterior wall of the maxillary sinus, we observed 88.89% sensitivity and 100% specificity.

Out of 84 fracture sites 14 false negative values of ultrasonography were observed. Among the 14 false negative values, six were fronto-zygomatic processes having 42.86% false value. Two of them were at the anterior/lateral wall of the maxillary sinus. Hence, at the fronto-zygomatic suture, we observed a 33.33% sensitivity of ultrasonography. We observed four false-positive values of ultrasonography of which the majority of them were located at the supraorbital margin with 92.10% at this site. The previous study of Jenkins [7] observed 86% sensitivity and 85% specificity of ultrasound for the diagnosis of orbital floor fracture. On contrary, our study failed to diagnose orbital floor fractures by using ultrasonography. We also observed that ultrasonography gave one false positive value at a mandibular angle and one false negative value at the mandibular condyle due to an undisplaced fracture of the neck. On contrary, 25 false negative values were observed in conventional radiographs. However, in our study, conventional radiographs reported zero false-positive value when compared to ultrasonography.

5. CONCLUSION

In conclusion, the current study depicts that ultrasonography is an economical, useful diagnostic tool for examining the bony fractures of facial trauma with a better sensitivity

rate when compared to conventional radiographs.

CONSENT AND ETHICAL APPROVAL

As per international standard or university standard guideline Patient's consent and ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. McCann PJ, Brocklebank LM, Ayoub AF. Assessment of zygomatico-orbital complex fractures using ultrasonography. *Br J Oral Maxillofac Surg.* 2000;38(5): 525-9.
DOI:
DOI:<https://doi.org/10.1054/bjom.2000.0501>
2. Friedrich RE, Plambeck K, Bartel-Friedrich S, Giese M, Schmelzle R. Limitations of B-scan ultrasound for diagnosing fractures of the mandibular condyle and ramus. *Clin Oral Investig* 2001;5(1):11-6.
3. Nezafati S, Javadrashid R, Rad S, Akrami S. Comparison of ultrasonography with submentovertex films and computed tomography scan in the diagnosis of zygomatic arch fractures. *Dentomaxillofacial Radiol* 2010;39(1): 116.
DOI:<https://doi.org/10.1259/dmfr/97056817>
4. Rama Mohan K, Koteswara Rao N, Leela Krishna G, Santosh Kumar V, Ranganath N, Vijaya Lakshmi U. Role of ultrasonography in oral and maxillofacial surgery: A review of literature. *J Maxillofac Oral Surg.* 2015;14(2):16270.
DOI:<https://doi.org/10.1007/s12663-014-0616-x>
5. Friedrich RE, Heiland M, Bartel-Friedrich S. Potentials of ultrasound in the diagnosis of midfacial fractures. *Clin Oral Investig.* 2003;7(4):226-9.
DOI:<https://doi.org/10.1007/s00784-003-0232-5> of orbital.
6. Jank S, Emshoff R, Etzelsdorfer M, Strobl H, Nicasi A, Norer B. Ultrasound versus computed tomography in the imaging floor fractures. *004 J Oral Maxillofac Surg.* 2004;62(2): 150-4.
7. Jenkins CN, Thuau H. Ultrasound imaging in assessment of fractures of the orbital floor. *Clin Radiol.* 1997;52(9):708-11.
DOI:[https://doi.org/10.1016/s0009-9260\(97\)80037-2](https://doi.org/10.1016/s0009-9260(97)80037-2)
8. zygomaticomaxillary Ogunmuyiwa SA, Fatusi OA, Ugboko VI, Ayoola OO, Maaji SM. The validity of ultrasonography in the diagnosis of complex fractures. *Int J Oral Maxillofac Surg* 2012;41(4): 500-5.
DOI:<https://doi.org/10.1016/j.ijom.2012.01.002>
9. Rajesh P, Rai AB Das. A comparison between radiography and ultrasonography in the diagnosis of zygomatic arch fracture. *Indian J Dent Res.* 2003;14(2):75-9.
10. Abu-Samra M, Selmi G, Mansy H, Agha M. Role of intra-operative ultrasound-guided reduction of nasal bone fracture in patient satisfaction and patient nasal profile (a randomized clinical trial). *Eur Arch Otorhinolaryngol.* 2011;268(4): 541-6.
DOI:<https://doi.org/10.1007/s00405-010-1401-1>
11. Javadrashid R, Khatoonabad M, Shams N, Esmaeili F, Jabbari Khamnei H. Comparison of ultrasonography with computed tomography in the diagnosis of nasal bone fractures. *Dentomaxillofacial Radiol.* 2011;40(8):486-91.
DOI:<https://doi.org/10.1259/dmfr/64452475>
12. Mohammadi A, Ghasemi-Rad M. Nasal bone fracture - Ultrasonography or computed tomography? *Med Ultrason.* 2011;13(4):292-5.
13. Gülicher D, Krimmel M, Reinert S. The role of intraoperative ultrasonography in zygomatic complex fracture repair. *Int J Oral Maxillofac Surg.* 2006;35(3):224-30.
DOI:<https://doi.org/10.1016/j.ijom.2005.10.005>
14. Lee JH, Park JH. The clinical usefulness of ultrasound-aided fixation using an absorbable plate system in patients with zygomatico-maxillary fracture. *Arch Plast Surg.* 2013;40(4):330-4.
DOI:<https://doi.org/10.5999/aps.2013.40.4.330>
15. Adeyemo WL, Akadiri OA. A systematic review of the diagnostic role of ultrasonography in maxillofacial fractures. *Int J Oral Maxillofac Surg.* 2011;40(7):655-61.

- DOI:<https://doi.org/10.1016/j.ijom.2011.02.001>
16. Rajeev A, Pai KL, Smriti K, Kadavigere R, et al. Diagnostic Accuracy of Ultrasonography in the Assessment of Facial Fractures. *Pesqui. Bras. Odontopediatria Clín. Integr.* 2019;19:e4832.

© 2021 Kamran and Al-Gorjia; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/78679>