

Journal of Geography, Environment and Earth Science International

22(4): 1-15, 2019; Article no.JGEESI.46598 ISSN: 2454-7352

Statistical Comparative Study of the Trend and Variation of Meteorological Parameters in Abeokuta, South-West Nigeria

M. A. Sodunke^{1*}, M. O. Sulaimon², R. S. Sunmonu¹, A. Mabosanyinje², Y. B. Lawal³, K. J. Oseni¹ and O. O. Oni⁴

¹Physics/Electronics Unit, Department of Science Laboratory Technology, Moshood Abiola Polytechnic, Abeokuta, Ogun, Nigeria. ²Department of Statistics and Mathematics, Moshood Abiola Polytechnic, Abeokuta, Ogun State, Nigeria.

³Department of Physical Sciences, University of Africa, Toru-orua, Bayelsa State, Nigeria. ⁴Department of Physics, Tai Solarin University of Education, Ijebu-Ode, Ogun State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author MAS was fully involved in the study design, data acquisition and drafting of the manuscript. Authors MOS and AM managed the statistical analyses. Authors RSS, YBL, KJO and OOO managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI:10.9734/JGEESI/2019/v22i430154 <u>Editor(s):</u> (1) Dr. Isidro Alberto Pérez Bartolomé, Professor, Department of Applied Physics, Faculty of Sciences, University of Valladolid, Spain. (2) Dr. Anthony R. Lupo, Professor, Department of Soil, Environmental, and Atmospheric Science, University of Missouri, Columbia, USA. (1) Uznir Ujang, Universiti Teknologi Malaysia, Malaysia. (2) Salvatore Grimaldi, Università degli studi della Tuscia, Italy. (3) Breno Barra, Federal University of Port Harcourt, Nigeria. (4) Nwaerema Peace, University of Port Harcourt, Nigeria. Complete Peer review History: <u>http://www.sdiarticle3.com/review-history/46598</u>

> Received 10 March 2019 Accepted 22 May 2019 Published 08 August 2019

Original Research Article

ABSTRACT

This study focuses on the statistical comparative study of the trend and variation of meteorological parameters covering a 10 year period (2001-2010) in the capital and largest city of Ogun State, Abeokuta, southwest region of Nigeria. The analyzed meteorological parameters were: wind speed, vapour pressure, relative humidity, temperature, sunshine and rainfall covering 10 years. The

*Corresponding author: E-mail: sodunke.mobolaji@mapoly.edu.ng;

calculated coefficient of variation (CV) for sunshine (22.78%), wind speed (21.55%), and rainfall (99.12%) is a proof of exceedance of variability of threshold of 10% while the CV calculated for air temperature (5.74%), relative humidity (4.52%) and vapour pressure (5.22%) show no significant variability. Significance test of meteorological parameters' trend reveals a notable reduction in the values of vapour pressure, air temperature and relative humidity. It is, however, difficult to argue for a well-defined change in most of the meteorological parameters based on the monthly time series analyses performed in this work. Only wind speed shows a statistically significant increasing trend during the period of observation at 1% significance level. The trend revealed by rainfall and sunshine is statistically not significant. ANOVA test of significant difference among meteorological parameters show a p-value (Sig.) of 0.000 is an indication of the significant difference in the analyzed mean monthly coefficient of variation for the meteorological parameters under study. The Tukey's multiple pair comparisons test, however, show that there is a significant difference between the mean monthly CV of rainfall-sunshine, rainfall-vapour pressure, rainfall-wind speed, rainfall-air temperature and rainfall-relative humidity. At the significance level of 5%, the calculated mean monthly CV of rainfall is significantly different from the mean monthly CV of other climatic parameters.

Keywords: Meteorological parameters; coefficient of variation; variability threshold; significance level; the significance of variability.

1. INTRODUCTION

Climate study is worth investigating since human lives are strictly attached and it is important to have knowledge of our environmental changes so that we would not be caught unaware by the consequences of the adverse effect offered. Climate change is associated with weather conditions and it has a great effect on environmental changes. Full knowledge of the environment will enhance proper management of risks so as to avert disasters. It is of great importance to know that improper management of climate could lead to natural disasters. It is therefore imperative to introduce protective schemes through the results obtained from quality research works that relates to environment. Year to year variability is caused by climate and has a link with socio-economic and environmental activities. It is of great importance toward the development and proper planning of schemes that relates to water resources such as the management of drought, the prevention and control of flood. Importantly, natural and ecosystems coupled with the society as a whole are directly linked to the consequences of changes in a climatic pattern either positively or negatively. Invariably, there could be alteration in the location of the major crop production regions on the earth due to climatic variability.

The variation of climate has a great influence on socio-economic activities. Research works conducted by [1] and [2] show that climatic parameters are closely related and have an influence on crop production. Precipitation, temperature, wind, pressure and humidity are physical conditions in the environment and atmosphere which are termed as weather because they have direct or indirect consequences upon the biosphere while the pattern of weather in a region over the period of time is referred to as climate [3]. The rise in the number of vehicles and industries are also factors in contemporary trends in climate [3]. It was reported by [4] that it is not only soil and pests that offers drawbacks in crop production but the effect of climate is of much influence.

Instability of weather could offer adverse effects in social, economic and regional competitiveness [5]. The negative change in the climatic pattern could be harmful to socio-economic activities thereby causing a reduction in food and fibers delivery to the teeming population [6]. The study conducted by [7] revealed that meteorological parameters from monthly series are of decreasing trend and not statistically significant except for rainfall and humidity that show an increasing trend which is statistically not significant.

The objectives of this study are to; examine the variations in rainfall, sunshine, air temperature, wind speed, relative humidity and water vapour patterns in the study area, examine the statistical link between sunshine, air temperature, wind speed, relative humidity rainfall and water vapour in the study area, determination of the trend of the meteorological parameters and presentation of their possible effects to the residents of the study area.




Fig. 1. The map of Nigeria showing the position of Abeokuta, Ogun State [8]

1.1 Study Area

Fig. 1 shows the study area (Abeokuta) lies between longitude and latitude of 4°19'20" E and 7°45'20" N respectively [8]. Abeokuta lies in the wooded savanna and the surface is characterized with masses of granite with grey colour. It covers an extensive area being surrounded by mud walls which are of 18 miles in extent [9].

3. METHODOLOGY

Ten years meteorological parameters (relative humidity, temperature, sunshine, wind speed, rainfall and vapour pressure) for Abeokuta Southwest Nigeria were collected from the Nigerian Meteorological Agency (NIMET) archive. The coefficient of variation was calculated as described in equation (1) by [10].

$$CV = \left(\frac{\sigma}{MP}\right) \times 100\% \tag{1}$$

The calculated monthly mean of the meteorological parameters is denoted as MP while σ is the standard deviation.

The statistical analyses were done using descriptive statistics, Kendall's tau_b, Spearman's rho, ANOVA and Tukey's multiple pair comparisons test. Data collected were analyzed electronically using Ms-Excel (version 2007) and SPSS (version 21.0).

The Kendall's tau_b for measuring order association between variables X and Y are given by the following formula:

$$\tau_b = \frac{P - Q}{\sqrt{D_r D_c}} \tag{2}$$

Where the P and Q listed above are double the "usual" P (number of concordant pairs) and Q (number of discordant pairs). Likewise, D_r is double the "usual" P+Q+X₀ (the number of concordant pairs, discordant pairs, and pairs on

which the row variable is tied) and D_c is double the "usual" P+Q+Y₀ (the number of concordant pairs, discordant pairs, and pairs on which the column variable is tied).

The Spearman correlation coefficient is defined as the Pearson correlation coefficient between the ranked variables.

For a sample of size *n*, the *n* raw scores X_{i} , Y_{i} are converted to ranks rg X_{i} , rg Y_{i} , and r_s is computed from:

$$r_{s} = \rho_{rgX,rgY} = \frac{Cov(rg_{X}, rg_{Y})}{\sigma_{rgX}\sigma_{rgY}}$$
(3)

Where ρ denotes the usual Pearson correlation coefficient but applied to the rank variables.

 $Cov(rg_X, rg_Y)$ is the covariance of the rank variables.

 σ_{rgX} and σ_{rgY} are the standard deviations of the rank variables.

Only if all n ranks are distinct integers, it can be computed using the popular formula

$$r_s = 1 - \frac{6\sum d_i^2}{n(n^2 - 1)}$$
(4)

Where $d_i = rg(X_i) - rg(Y_i)$, is the difference between the two ranks of each observation. *n* is the number of observations.

ANOVA (One-way) is a technique that can be used to compare means of two or more samples (using the F distribution). The ANOVA test the

4. RESULTS

• Data presentation

null hypothesis that samples in all groups are drawn from populations with the same mean values.

The normal linear model implemented in this study is means model which is given as :

$$y_{ij} = \mu_j + \epsilon_{ij} \tag{5}$$

Tukey's multiple pair comparisons test compares the means of every treatment to the means of every other treatment; that is, it applies simultaneously to the set of all pairwise comparisons $\mu_i - \mu_j$ and identifies any difference between two means that is greater than the expected standard error.

The formula for Tukey's test is:


$$q_s = \frac{Y_A - Y_B}{SE} \tag{6}$$


Where Y_A is the larger of the two means being compared.


 Y_B is the smaller of the two means being compared.


SE is the standard error of the two sums of the means.

This q_s value can then be compared to a q value from the studentized range distribution. If the q_s value is larger than the critical value q_{α} obtained from the distribution, the two means are said to be significantly different at level α , $0 \le \alpha \le 1$.










Sodunkeet al.; JGEESI, 22(4): 1-15, 2019; Article no.JGEESI.46598










Fig. 2. Monthly mean values and trend of the meteorological parameters in abeokuta




Sodunkeet al.; JGEESI, 22(4): 1-15, 2019; Article no.JGEESI.46598












Fig. 3. Monthly coefficient of variation (cv) and trend of the meteorological parameters in abeokuta




Fig. 4. Time series plot of observed &estimated (Trend Value) CV

• Data analysis

Table 1. Descriptive statistics of monthly coefficient of variation (CV)

	N	Minimum	Maximum	Mean	Std. Deviation
Rainfall	12	25.62	283.23	99.1247	83.36912
Sunshine	12	8.01	41.84	22.7808	11.35960
Vapour pressure	12	1.41	17.33	5.2167	5.36420
Wind speed	12	18.04	28.95	21.5492	3.12828
Air temperature	12	5.00	7.13	5.7400	.57479
Relative humidity	12	1.16	14.63	4.5150	4.74239
Valid n (listwise)	12				

Table 2. Bivariate Correlations Among Meteorological Parameters

			Rainfall	Sunshine	Vapour pressure	Wind speed	Air temperature	Relative humidity
Kendall's	Rainfall	Correlation Coefficient	1.000	.273	.545 [*]	.091	121	.545 [*]
tau_b		Sig. (2-tailed)		.217	.014	.681	.583	.014
		Ν	12	12	12	12	12	12
	Sunshine	Correlation Coefficient	.273	1.000	.121	.030	242	061
		Sig. (2-tailed)	.217		.583	.891	.273	.784
		Ν	12	12	12	12	12	12
	Vapour	Correlation Coefficient	.545*	.121	1.000	.182	152	.636**
	pressure	Sig. (2-tailed)	.014	.583		.411	.493	.004
		N	12	12	12	12	12	12
	Wind speed	Correlation Coefficient	.091	.030	.182	1.000	.000	121
		Sig. (2-tailed)	.681	.891	.411		1.000	.583
		N	12	12	12	12	12	12
	Air temperature	Correlation Coefficient	121	242	152	.000	1.000	091
		Sig. (2-tailed)	.583	.273	.493	1.000		.681
		N	12	12	12	12	12	12
	Relativehumidity	Correlation Coefficient	.545*	061	.636**	121	091	1.000
	-	Sig. (2-tailed)	.014	.784	.004	.583	.681	
		N	12	12	12	12	12	12

Sodunkeet al : IGEESI	$22(A) \cdot 1_{-}15$	2019; Article no.JGEESI.46598
Souurikeel al., JGEESI	22(4). 1-10,	2019, AILICIE 110.JGEE31.40390

			Rainfall	Sunshine	Vapour pressure	Wind speed	Air temperature	Relative humidity
Spearman's	Rainfall	Correlation Coefficient	1.000	.315	.748	.140	252	.706
ho		Sig. (2-tailed)		.319	.005	.665	.430	.010
		N	12	12	12	12	12	12
	Sunshine	Correlation Coefficient	.315	1.000	.126	.049	322	273
		Sig. (2-tailed)	.319		.697	.880	.308	.391
		N	12	12	12	12	12	12
	Vapour	Correlation Coefficient	.748 ^{**}	.126	1.000	.238	126	.762**
	pressure	Sig. (2-tailed)	.005	.697		.457	.697	.004
		N	12	12	12	12	12	12
	Wind speed	Correlation Coefficient	.140	.049	.238	1.000	.021	112
		Sig. (2-tailed)	.665	.880	.457		.948	.729
		N	12	12	12	12	12	12
	Air temperature	Correlation Coefficient	252	322	126	.021	1.000	119
		Sig. (2-tailed)	.430	.308	.697	.948		.713
		N	12	12	12	12	12	12
	Relative	Correlation Coefficient	.706 [*]	273	.762**	112	119	1.000
	humidity	Sig. (2-tailed)	.010	.391	.004	.729	.713	
	-	Ň	12	12	12	12	12	12

*Correlation is significant at the 0.05 level (2-tailed)**Correlation is significant at the 0.01 level (2-tailed)

Table 3: Significance test of meteorological parameters' trend significance levels are Indicated: 95% (*), 99% (**)

Meteorological Parameter	Kendall's tau_b	Spearman's rho	Pearson
Rainfall	0.091	0.091	0.061
Sunshine	-0.030	0.028	0.065
Vapour pressure	-0.182	-0.245	-0.380
Wind speed	0.576** (increasing trend)	0.713** (increasing trend)	0.712** (increasing trend)
Air temperature	-0.121	-0.133	0.014
Relative humidity	-0.303	-0.343	-0.460

Sodunkeet al.; JGEESI, 22(4): 1-15, 2019; Article no.JGEESI.46598

Table 4. ANOVA Test of significant difference among meteorological parameters

	Sum of Squares	df	Mean Square	F	Sig.	
Between Groups	80160.369	5	16032.074	13.471	.000	
Within Groups	78546.594	66	1190.100			
Total	158706.963	71				

Table 5. Tukey's multiple pair comparisons test

(I) Group	(J) Group	Mean difference (I-J)	Std. Error	Sig.	95% Confidence interval	
					Lower bound	Upper bound
Rainfall	Sunshine	76.34417 [*]	14.08368	.000	35.0072	117.6811
	Vapour pressure	93.90833 [*]	14.08368	.000	52.5714	135.2453
	Wind speed	77.57583 [*]	14.08368	.000	36.2389	118.9128
	Air temperature	93.38500 [*]	14.08368	.000	52.0481	134.7219
	Relative humidity	94.61000 [*]	14.08368	.000	53.2731	135.9469
Sunshine	Rainfall	-76.34417*	14.08368	.000	-117.6811	-35.0072
	Vapour pressure	17.56417	14.08368	.812	-23.7728	58.9011
	Wind speed	1.23167	14.08368	1.000	-40.1053	42.5686
	Air temperature	17.04083	14.08368	.830	-24.2961	58.3778
	Relative humidity	18.26583	14.08368	.786	-23.0711	59.6028
Vapour pressure	Rainfall	-93.90833 [*]	14.08368	.000	-135.2453	-52.5714
	Sunshine	-17.56417	14.08368	.812	-58.9011	23.7728
	Wind speed	-16.33250	14.08368	.854	-57.6694	25.0044
	Air temperature	52333	14.08368	1.000	-41.8603	40.8136
	Relative humidity	.70167	14.08368	1.000	-40.6353	42.0386
Wind speed	Rainfall	-77.57583 [*]	14.08368	.000	-118.9128	-36.2389
	Sunshine	-1.23167	14.08368	1.000	-42.5686	40.1053
	Vapour pressure	16.33250	14.08368	.854	-25.0044	57.6694
	Air temperature	15.80917	14.08368	.870	-25.5278	57.1461
	Relative humidity	17.03417	14.08368	.831	-24.3028	58.3711

	00(4) 4 45	
Sodunkeet al.; JGEESI,	22(4): 1-15	, 2019; Article no.JGEESI.46598

(I) Group	(J) Group	Mean difference (I-J)	Std. Error	Sig.	95% Confidence interval	
					Lower bound	Upper bound
Air temperature	Rainfall	-93.38500 [*]	14.08368	.000	-134.7219	-52.0481
	Sunshine	-17.04083	14.08368	.830	-58.3778	24.2961
	Vapour pressure	.52333	14.08368	1.000	-40.8136	41.8603
	Wind speed	-15.80917	14.08368	.870	-57.1461	25.5278
	Relative humidity	1.22500	14.08368	1.000	-40.1119	42.5619
Relative humidity	Rainfall	-94.61000 [*]	14.08368	.000	-135.9469	-53.2731
-	Sunshine	-18.26583	14.08368	.786	-59.6028	23.0711
	Vapour pressure	70167	14.08368	1.000	-42.0386	40.6353
	Wind speed	-17.03417	14.08368	.831	-58.3711	24.3028
	Air temperature	-1.22500	14.08368	1.000	-42.5619	40.1119

*. The mean difference is significant at the 0.05 level

Group	N	Subset for alpha = 0.05		
		1	2	
Relative humidity	12	4.5150		
Vapour pressure	12	5.2167		
Air temperature	12	5.7400		
Wind speed	12	21.5492		
Sunshine	12	22.7808		
Rainfall	12		99.1250	
Sig.		.786	1.000	

Table 6. Tukey's homogeneous subsets

Means for groups in homogeneous subsets are displayed a. Uses Harmonic Mean Sample Size = 12.000

5. DISCUSSION OF RESULTS

The mean monthly values and trend of the meteorological parameters in the study area are presented in Fig. 2. For sunshine, months of January, February, March, April, May, October, November and December show maximum sunshine regimes. Incessant cloud formation depletes the amount of sun reaching us is accountable for the minimum sunshine experienced in August which was earlier reported by [5] for Ibadan sunshine hour in 2012. R^2 of 0.011 implies that approximately 1.1% of the variation in sunshine distribution is being explained by the monthly period of study.

A gradual pick up in vapour pressure was experienced in January which spans through the months of February, March and April but dropped in May. There was persistence in the trend of vapour pressure in July and August. October and November showed a rise in the vapour pressure while there was a sudden collapse in December. R^2 of 4E-05 implies that the monthly period of study does not explain a significant variation in vapour pressure distribution.

Unstable wind speed distribution was observed for the period under study. February and March relay peak levels of wind speed while September marked a low distribution. R^2 of 0.639 implies that approximately 63.9% of the variation in wind speed distribution is being explained by the monthly period of study.

The highest value of air temperature was recorded in July which eventually collapsed in August and spans through August and September. Least temperature is shown in September which gradually increases from the months of October to December. R^2 of 0.003 implies that approximately 0.3% of the variation in air temperature distribution is being explained by the monthly period of study.

There is an exponential rise in relative humidity in months January, February, March, April, May, June and July. August revealed an upward trend movement of the relative humidity regimes which later showed a trend collapse and decreased from September to December. R^2 of 0.258 implies that approximately 25.8% of the variation in relative humidity distribution is being explained by the monthly period of study.

Mean monthly distribution of rainfall showed a low rainfall for the months of January, February, March, November and December. There was an upward increase in rainfall from April till July. A sudden collapse in the rising level of rainfall was experienced in August which later increased gradually in September and dropped down from the month of October to December. R^2 of 0.033 implies that approximately 3.3% of the variation in rainfall distribution is being explained by the monthly period of study.

Figs. 3 & 4 shows the coefficients of variation (CV) for the meteorological parameters under study. The highest values of the CV calculated for the climatic parameters sunshine, vapour pressure, wind speed, rainfall, air temperature and relative humidity were: 41.84%, 17.33%, 28.95%, 283.23%, 7.13% and 14.63% while lowest values were: 8.01%, 1.41%, 18.04%, 25.62%, 5.00% and 1.16% respectively. From the obtained results, rainfall exhibits the highest variation while relative humidity depicts the least variation.

The descriptive statistics result from Table 1, indicates that we expect the monthly CV for rainfall to be 99.12%, the expected monthly CV for sunshine to be 22.78%, the expected monthly CV for vapour pressure to be 5.22%, the expected monthly CV for wind speed to be 21.55%, the expected monthly CV for air temperature to be 5.74% and the expected monthly CV for relative humidity to be 4.52%.

Table 2 shows bivariate correlations among the meteorological parameters using Kendall's tau b and Spearman's rho statistics. From Kendall's tau b analysis, it shows there is a weak positive association between rainfall-sunshine, rainfall-Sunshine-vapour pressure, wind speed. sunshine-wind speed, vapour pressure-wind speed. The weak negative association is observed between rainfall-air temperature, wind speed-relative humidity, vapour pressure-air temperature, sunshine-air temperature and sunshine-relative humidity. There is an average significant relationship between rainfall-vapour pressure, rainfall-relative humidity but a strong positive significant correlation between vapour pressure and relative humidity at 5% significance level. There is no association between the air temperature-wind speed. Spearman's rho results show that there is a very strong positive significant correlation between rainfall-vapour rainfall-relative humidity, vapour pressure. pressure-relative humidity at 5% significance level. A negative correlation is experienced between rainfall-air temperature, sunshine-air temperature, sunshine-relative humidity, vapour pressure-air temperature, wind speed-relative humidity and air temperature-relative humidity.

Significance test of meteorological parameters' trend from Table 3 reveals a notable deterioration in the values of vapour pressure, air temperature and relative humidity. However, it is difficult to argue for a well-defined change in most of the meteorological parameters based on the monthly time series analysis performed in this work. Only wind speed shows a statistically significant increasing trend during the period of observation at 1% significance level. The trend shows by others are statistically not significant.

ANOVA Test of significant difference among meteorological parameters from Table 4 shows a p-value (Sig.) of 0.000 indicating a significant difference in the mean monthly coefficient of variation of the six climatic parameters (rainfall, sunshine, vapour pressure, wind speed, air temperature and relative humidity). In other words, the mean monthly coefficient of variation of at least one of the parameters is significantly different from others.

The Tukey's multiple pair comparisons test from Table 5 shows that there is a significant difference between the mean monthly CV of rainfall–sunshine, rainfall-vapour pressure, rainfall-wind speed, rainfall-air temperature and rainfall-relative humidity. It is therefore evident that the mean monthly CV of rainfall is significantly different from the mean monthly CV of the other climatic parameters at 5% significance level.

Tukey's homogeneous subset from Table 6 shows the order of importance of the meteorological parameters under study. It reveals that relative humidity is of the most important, followed by vapour pressure, air temperature, wind speed, sunshine and rainfall as the least important.

6. CONCLUSION

This study revealed the occurrence of a significant difference in variation for all the investigated meteorological parameters. Also, there is a notable deterioration in the values of vapour pressure, air temperature and relative humidity. Only wind speed shows a statistically significant increasing trend during the period of observation while the trend shown by others are statistically not significant. Rainfall, wind speed and temperature show tolerable values which are not life-threatening to the residents of Abeokuta. For vapour pressure, only months of January, August and December are safe but proper precautionary measures must be infused in other months in order to reduce problems of high blood pressure due to high vapour pressure. Since the relative humidity is higher than the tolerable limit of 60%, Abeokuta is subject to heat. Therefore. the residents should endeavor to provide themselves cooling systems for homeostasis to be engaged. Also, exposure time to sunlight should also be reduced in months of November and December respectively so as to avoid or reduce the ageing of the skin due to excessive exposure to sunlight.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- 1. Ayoade JO. Introduction to climatology for the Tropics, Revised edition Spectrum books limited, Ibadan; 2004.
- Cicek I, Turkogu N. Urban effects on precipitation Ankara Turkey. Atmosfera. 2005;18(3):173-187.
- Cunningham WP, Cunnigham MA, Sanjo B. A global concern. 8th ed, New York, published by McGraw-Hill. 2005;16-316.

- Efe SI. Climate change and Food Security in Africa, Delta State Nigeria. Conference proceedings on climate change and the Nigerian Environment held at UNN 29th June-July 2nd. 2009;105-126.
- Ogolo EO, Adeyemi B. Variations and trends of some metrological parameters at Ibadan, Nigeria. Pacific Journal. 2009; 10(2);981-987.
- Reddy KR, Hodges HF. Climate change and global crop productivity. CAB International Publishing. 2000;1-5.
- Syed ALS, Sobla N, Abdullah K, Rahman ZU, Syed MAS. Trends and variability in climate parameters of Peshawar district. Journal of Science Technology and Development. 2012;31(4):341-357.
- David OB, Lobina GP. Modelling a critical infrastructure-driven spatial database for proactive disaster management: A developing country context. Journal of Disaster Risk Studies. 2016;8(1):1-14.
- Sodunke MA, Adewale AO, Alabi AA, Sunmonu RS, Mabosanyinje A. Analytical study on the variation of climatic parameters at Abeokuta, south-west, Nigeria. International Journal of Advances in Science Engineering and Technology. 2016;4(4):25-26.
- Murat Turkes, Utku M. Sumer, Gonul Kilic. Variations and trends in annual mean air temperatures in Turkey with respect to climatic variability. International Journal of Climatology. 1995;15:557-564.

© 2019 Sodunke et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: http://www.sdiarticle3.com/review-history/46598