



## **A Case Report on Malignant Ovarian Germ Cell Tumor**

**Sanket V. Madavi<sup>a\*#</sup>, Samruddhi Gujar<sup>a</sup>, Ranjana Sharma<sup>a≡</sup>, Ruchira Ankar<sup>a</sup>,  
Savita Pohekar<sup>a</sup>, Sheetal Sakharkar<sup>a</sup>, Vaishali Tembhare<sup>a</sup>  
and Kanchan Bokade<sup>a</sup>**

<sup>a</sup> Department of Medical Surgical Nursing, Smt. Radhikabai Meghe Memorial College of Nursing,  
Datta Meghe Institute of Medical Sciences (Deemed To Be University) Sawangi (Meghe), Wardha,  
India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i54A33717

#### Editor(s):

(1) Dr. Paola Angelini, University of Perugia, Italy.

#### Reviewers:

(1) A. Anand Kumar, Sri Venkateswara Veterinary University, India.

(2) Sapna Singh, Narayan Nursing College, Gopal Narayan Singh University, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:  
<https://www.sdiarticle5.com/review-history/75004>

**Case Study**

**Received 01 September 2021**

**Accepted 05 November 2021**

**Published 08 December 2021**

### **ABSTRACT**

**Introduction:** A germ cell is a type of Tumor that arises from germ cell (GCT). Germ cell tumors are both cancerous and benign. The gonads contains majority of germ cell (ovary and testis). GCT that arise outside the gonads could be the result of embryo development errors.

**Clinical Findings:** Abdominal mass without pain, backache, and weight loss.

**Diagnostic Evaluation:** HCG and alphafetoprotein level in the blood identifying women with malignant ovarian germ cell tumor is beneficial. Serum alphafetoprotein and HCG is elevated as condition progress.

**Therapeutic Intervention:** Patient treated with inj. Bleomycin, inj. Cisplatin, inj. Etoposide, tab Pan 40 miligram and inj. Emset.

**Conclusion:** A 3year old female was admitted in AVBRH Sawangi Meghe wardha with the major complaint of right abdominal swelling for 6 month, both when they observed abdominal mass without pain and swelling was small in size and backache and weight loss.

<sup>#</sup> Basic B. Sc. Nursing Third Year Student;

<sup>≡</sup> Dr.;

\*Corresponding author: E-mail: sanketmadavi1309@gmail.com;

**Keywords:** Cancerous and benign germ-cell tumour; neoplasm; gonads; malignant.

## 1. INTRODUCTION

In terms of biology and clinical presentation and Juvenile GCTs are very similar to adult testicular GCT however there are few critical changes of aware of GCT incidents has two peak one in young children (0-4 aged) and other in puberty according to epidemiologic data [1].

While GCTs in adolescents and adults appear to have the same histology and molecular biology, germ cell tumours in very young children have substantial difference suggesting that maybe unique disease [2]. GCT roughly accounts for 3% malignancies in children aged 0–18 and their prevalence increases as they approach puberty GCTs account for 15% of all malignancies in children under the age of 18. Pediatric GCTs are treated with cisplatin based multiagent regimens, which have proven to be highly efficacious even in the setting of advanced disease, similar to adult GCTs [3].

### 1.1 Patient Identification

A 3 years old female with known case of malignant ovarian germ cell tumour come to the AVBRH Sawangi (Meghe), Wardha with Chief complaint of swelling on right abdominal region for 6 months both when they noticed abdominal mass without pain as well as swelling was small in size and backache, weight loss.

#### 1.1.1 Present medical history

A 3 years old female was admitted in AVBRH in paediatric ward on date 19/05/2021 with complaint of chief complaint of swelling on right abdominal region for 6 months both when they noticed abdominal mass without pain as well as swelling was small in size and backache, weight loss.

#### 1.1.2 Past medical history

Previously, the patient was treated in a Yavatmal civil hospital. There is no other medical history of hypertension, diabetes, asthma, Tuberculosis, or seizures in the patient.

#### 1.1.3 Family history

My patient has 3 family members including father, mother and herself. Her parents are

healthy and has no relevant history of genetic abnormalities contributing to patient's condition.

#### 1.1.4 Past interventions and outcome

The patient was admitted in civil hospital Yavatmal. Patient's general conditions was poor; hence, she was referred to AVBRH Sawangi (Meghe), Wardha for further management.

## 1.2 Clinical Findings

Abdominal mass without pain, backache, and weight loss.

### 1.2.1 Aetiology

This slow-growing malignancy starts in female germ cells and leads to a disruption in the hormone beta-human chorionic gonadotropin (beta-hCG), which the body generates in greater amounts during pregnancy. The most prevalent kind of ovarian germ cell tumour is dysgerminoma [4].

### 1.2.2 Physical examination

A 3 years old female look malnourished and had abdominal mass without pain along with backache and weight loss.

### 1.2.3 Diagnostic assessment

Physical review on the basis of patient history, physical examination and other all blood investigation and biopsy done. Blood test: Hb= 11.1gm%, RBC= 4.5, WBC= 7000, PLT= 2.74l lakhs/cu mm, CRP=0.50, Bilirubin= 0.4, Total protein=6.6, INR = 1.07, HCG= 10 mIU/ml. Tumour markers: AFP= 400 ng/ml. Urine: Albumin and sugar normal. Ultrasonography: Right ovarian mass.

## 2. MANAGEMENT

### 2.1 Medical Management

Inj. Bleomycin (10 to 20 unit/m<sup>2</sup>), Inj. Cisplatin (20 miligram/m<sup>2</sup>), inj. Etoposide (50 miligram/m<sup>2</sup>), Emset (2 milliliter), tab Pan 40 mg.

## 2.2 Surgical Management

### 2.2.1 Nursing management

Observed for side effect. Monitored WBC and platelet count prior to each dose. Observed for signs of allergic reaction. Taught patient's parents about signs of bleeding.

### 2.2.2 Follow-up care

Because around 75% of MOGCT recurrences happen during the first year, vigorous early follow-up every 4-8 weeks is reasonable and should be audited 229 The peritoneal cavity and, less commonly, the retroperitoneal lymph nodes or lungs are common recurrence locations [5]. The goals of follow-up should be to assess therapy response, manage treatment-related problems, and discover persistent or recurring disease early [6]. Tumor markers should be assessed at each appointment, and chest X-rays should be done every 2 months for the first 2 years, then every 3-6 months for the next 3 years. Women should be evaluated for unfavourable physical and cardiovascular effects, as well as psychosocial and psychosexual repercussions of treatment, in addition to disease-related follow-up [7].

## 3. DISCUSSION

A 3 years old female was admitted in AVBRH Sawangi (Meghe), Wardha with chief complaint of swelling on right gluteal region, fever, weakness, irritability, constipation, chest pain, unusual shape and size for further management [8]. After all investigation the Doctor diagnosed case as malignant ovarian Germ cell tumour .after treatment patient condition was stable [9]. Malignant ovarian Germ cell tumour is a clinical disorder, which is commonly seen in patients. Germ cell cancers are caused by molecular abnormalities in primordial germ cells, which are early germline progenitors [10].

These cells go from the yolk sac to the gonadal ridge's midline in the developing embryo [11-13]. Extragonadal GCTs have a preference towards the midline since the migration mechanism has been disrupted [14]. The pluripotent nature of these cells explains the differences in histology reported [15]. Immature teratomas have less differentiated tissue, such as neuroectoderm, and are categorised (I-3) according to their dedifferentiation. Despite the fact that both

mature and immature teratomas are benign tumours, immature teratomas, particularly those with high grade characteristics, require a careful histologic examination due to their documented relationship with microscopic foci of malignancy [16].

## 4. CONCLUSION

In postmenopausal women, malignant ovarian germ cell tumours are an extremely rare occurrence [17]. Malignant mixed germ cell tumours of the ovary are a highly aggressive neoplasm, and any adolescent female presenting with a rapidly increasing pelvic mass requires early care and fertility sparing surgery [18,19,20]. This series demonstrates the excellent prognosis for females with ovarian germ cell tumours, with a recurrence rate of 4.5 percent and a mortality rate of 3%. We stress the necessity of immunohistochemical investigation and serum AFP measurement in determining the accurate diagnosis, and we emphasise the likelihood of endometriosis or malignancies resulting from endometriosis as precursor causes. These people have a poor prognosis, regardless of whether they present with early-stage illness or whether an epithelial component is discovered [21]. Adjuvant therapy with platinum-based medicines is recommended, but more research is needed to understand why this disease has such a poor prognosis in this age group [22].

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

It is not applicable.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Frazier AL, Amatruda JF. Germ cell tumors. In: Fisher DE, Nathan D, Look AT., editors. Nathan and Oski's Textbook of Pediatric Hematology-Oncology. London, UK: Elsevier; 2009. [Google Scholar]
2. Huddart RA, Norman A, Shahidi M, et al. Cardiovascular disease as a long-term

- complication of treatment for testicular cancer. *Journal of Clinical Oncology*. 2003;21(8):1513–1523.  
DOI: 10.1200/jco.2003.04.173. [PubMed] [CrossRef] [Google Scholar]
3. Travis LB, Curtis RE, Storm H, et al. Risk of second malignant neoplasms among long-term survivors of testicular cancer. *Journal of the National Cancer Institute*. 1997;89(19):1429–1439.  
DOI: 10.1093/jnci/89.19.1429. [PubMed] [CrossRef] [Google Scholar]
  4. Travis LB, Fossa SD, Schonfeld SJ, et al. Second cancers among 40,576 testicular cancer patients: focus on long-term survivors. *Journal of the National Cancer Institute*. 2005;97(18):1354–1365.  
DOI: 10.1093/jnci/dji278. [PubMed] [CrossRef] [Google Scholar].
  5. David M Gershenson *Cancer*. 1993;71(S4):1581-1590,.
  6. Christine A Parkinson, Helen M Hatcher, Thankamma V Ajithkumar. *Obstetrical & gynecological survey*. 2011;66(8):507-514,.
  7. David M Gershenson. *Journal of Clinical Oncology*. 2007;25(20):2938-2943.
  8. Solheim, J Kærn, Claes G Tropé, E Rokkones, AA Dahl, JM Nesland, SD Fosså *Gynecologic oncology*. 2013;131(2) :330-335.
  9. Zanagnolo V, Sartori E, Galleri G, Pasinetti B, Bianchi U. *European Journal of Gynaecological oncology*. 2004;25(3):315-320,
  10. Chow SN, Yang JH, Lin YH, Chen YP, Lai JI, Chen RJ, Chen CD. *International Journal of Gynecology & Obstetrics*. 1996;53(2):151-158.
  11. Jeong-Yeol Park, Dae-Yeon Kim, Dae-Shik Suh, Jong-Hyeok Kim, Yong-Man Kim, Young-Tak Kim, Joo-Hyun Nam *Annals of surgical oncology* 2016;23 (9):2982- 2987.
  12. Zhang R, Sun YC, Zhang GY, Wu LY, Zuo J. *European Journal of Gynaecological Oncology*. 2012;33 (5):489-492.
  13. Antoine De Backer, Gerard C Madern, J Wolter Oosterhuis, Friederike GAJ Hakvoort-Cammel, Frans WJ Hazebroek *Pediatric blood & cancer*. 2006;46(4):459-464.
  14. Lori Hoffner, Susan Shen-Schwarz, Ranjan Deka, Aravinda Chakravarti, Urvashi Surti *Cancer genetics and cytogenetics*. 1992;62(1):58-65.
  15. Jin Li, Xiaohua Wu *Current treatment options in oncology*. 2016;17 (8):1-12.
  16. Stergios Boussios, Ayoma Attygalle, Steve Hazell, Michele Moschetta, Jennifer Mclachlan, Alicia Okines, Susana Banerjee. *Anticancer research* 35 (12), 6713-6722, 2015.
  17. 150. Ghavat, Chinmay, Nitin Bhola, Anendd Jadhav, Nupoor Deshpande, and Chetan Gupta. Malignant transformation of solitary cylindroma involving the frontonasal region- A rarity. *Journal of Clinical and Diagnostic Research*. 2020; 14(4):ZD1–4.  
Available:<https://doi.org/10.7860/JCDR/2020/43006.13620>
  18. Goyal Richa, Chandra Veer Singh, Shraddha Jain, Sana Parveen, Prasheelkumar Gupta, and Deepak Gupta. Our experience with nasolabial flaps in soft tissue reconstruction of oral malignancy; Feasible option in rural set up. *Indian Journal of Otolaryngology and Head & Neck Surgery*, n.d.  
Available:<https://doi.org/10.1007/s12070-020-02257-2>
  19. Shelke Uttara Vijay, Shourya Acharya, Deepti Sandeep Shrivastava. Co-existence of chronic lymphocytic leukaemia and malignancy of uterine cervix. *Journal of Evolution of Medical and Dental Sciences-JEMDS*. 2020; 9(18):1522–24.  
Available:<https://doi.org/10.14260/jemds/2020/332>
  20. Toshniwal Shikha M, Saunitra A Inamdar, Himanshi D Agarwal, Sakshi R Sharma. Malignant brenner tumor- A rare case of ovarian carcinoma. *Medical Science*. 2020;24(105):3550–54.
  21. Gawri DJ, Agrawal DM, Waigi R. A comparative study of laparoscopic assisted vaginal hysterectomy vs nondescent vaginal hysterectomy for benign gynaecological diseases. *International Journal of Pharmaceutical Research*. 2019;11:2018–2021.  
Available:<https://doi.org/10.31838/ijpr/2019.11.04.503>

22. Jaiswal N, Makrande J, Bhake A, Bapat AV. Ema, vimentin, desmin, calretinin, e-cadherin on cell block to differentaite adenocarcinoma cells from benign reactive mesothelial cells. International Journal of Pharmaceutical Research. 2019;11:1860–1864.  
Available:<https://doi.org/10.31838/ijpr/2019.11.02.213>

---

© 2021 Madavi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/75004>*