

Population Structure and Regeneration of *Croton scabiosus* Bedd. (Euphorbiaceae) in Dry Forests of South Eastern Ghats of YSR Kadapa District, Andhra Pradesh, India

L. Nagireddy¹, S. Nazaneen Parveen¹ and T. Pullaiah^{2*}

¹Department of Environmental Science, Yogi Vemana University, Kadapa-516003, Andhra Pradesh, India.

²Department of Botany, Sri Krishnadevaraya University, Anantapuramu-515003, Andhra Pradesh, India.

Authors' contributions

This work was carried out in collaboration between all authors. Author TP designed the study and wrote the protocol. Author LN carried out the field study and wrote the first draft of the manuscript. Author SNP supervised the work, did statistical analysis, corrected the first draft and managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2017/31498

Editor(s):

- (1) Renata Guimarães Moreira-Whitton, Departamento de Fisiologia - Instituto de Biociências-USP, Cidade Universitária, Brazil.
(2) Aleksander Lisowski, Warsaw University of Life Sciences, Department Agricultural and Forestry Engineering, Poland.

Reviewers:

- (1) Martin Potgieter, University of Limpopo, South Africa.
(2) Surendra Singh Bargali, Kumaun University, Nainital, India.
(3) Özlem Sultan Aslantürk, Adnan Menderes University, Turkey.

Complete Peer review History: <http://www.sciencedomain.org/review-history/18693>

Original Research Article

Received 9th January 2017
Accepted 7th April 2017
Published 19th April 2017

ABSTRACT

Croton scabiosus Bedd. (Euphorbiaceae), an endemic tree of dry forests of South Eastern Ghats, is categorized a vulnerable tree. For effective conservation of this species studies on its population structure and regeneration are required. The population structure and regeneration of *Croton scabiosus* were studied in 11 localities of dry forests of Kadapa district, India, by laying 66 quadrats 5x200 m in size. A total of 2571 individual trees with a range of 61-100 individuals in each quadrat were recorded. The majority (81%) of the individuals are in 10-20 cm and 21-30 cm gbh class,

*Corresponding author: E-mail: pullaiah.thammineni@gmail.com;

followed by 19% of 31-40 cm gbh class. The height class distribution revealed that 74.06% trees are in 2-3 m height class, while 20.78% trees in 4-5 m height class and only 5.14% trees are 6-8 m height. In the regeneration category 8.8% of seedlings and 49.7% saplings were recorded. In addition, *C. scabiosus* is associated with 54 tree species belonging to 39 genera and 21 families. The most commonly associated tree species are; *Acacia caesia*, *A. chundra*, *Cleistanthus collinus*, *Gardenia gummifera*, *Mimosa intsia*, *Ochna obtusa*, *Strychnus nux-vomica*, *Pterocarpus santalinus* and *Wrightia tinctoria*. In certain sites *C. scabiosus* formed the dominant lower canopy tree in the elevation range of 155 m-328 m.

Keywords: *Croton scabiosus*; endemic tree; vulnerable; dry forests; associated trees; population structure; regeneration.

1. INTRODUCTION

Tropical dry forest accounts for 38.2% of the total forest cover of India [1]. These dry forests not only constitute rich species diversity, generate a variety of natural resources, but also promote livelihood security to local communities [2]. Tropical dry forests occur in climate with seasonal rainfall, with a mean annual rain fall of 250-2000 mm and Potential Evapotranspiration (PET) of >1 [3]. The distribution of plants is determined by climatic variables like rain fall, temperature, soil conditions, moisture, nutrient availability and competition between plant species for crown and root space as well as human interference [4].

Croton scabiosus Bedd. (Euphorbiaceae), a small deciduous tree of 4-6 m in height, is endemic to South Eastern Ghats of Andhra Pradesh [5-7]. This species is represented by small populations with a scattered distribution in dry deciduous forests (Figs. 1, 2) of South Eastern Ghats of Andhra Pradesh [8]. It is principally found in Ananthapuramu, Kadapa and Nellore Districts of Southern Andhra Pradesh, India. It is listed as vulnerable according to IUCN Red list status [8]. The species is medicinally important, as the seeds are used as an antidote for snake bite and scorpion sting by the local people [9]. It has a very attractive silvery to golden foliage, which can be recommended as an ornamental garden plant [10]. Its populations in the natural habitat in Kadapa forests are primarily threatened by forest fires, fungal infection of seeds and poor germination [7]. As this is an endemic and medicinally important species and its populations are small and declining in the native habitats, studies on regeneration and phytosociological information are necessary for its conservation.

2. MATERIALS AND METHODS

2.1 Study Area

YSR Kadapa district is located between 13°43' and 15°14' North latitude and 77°55' and 79°29' East longitude. The total geographical area of the district is 15,379 km², with the forest area comprising 5,050 km², which is 32.87% of the district area. Day temperature reaches maximum 46°C in May and minimum 31°C in December in Kadapa district. Average rainfall is 700 mm. The present study was undertaken in the dry deciduous forests of Kadapa district in 11 places namely; Idupulapaya, Palakondalu, Nithya pooja Kona, Porumamilla, Joythi, Kanampalli (Pulivendula), Sundupalli, Sanipaya, Polathala, Guvvalacheruvu, and Chitvel-Rapur hills. These 11 locations are identified based on forest vegetation of *C. scabiosus* species distribution (Figs. 1 and 2). The study was carried out at each location to assess the density 4.04/0.6 ha⁻¹ in totally 6.6ha population and regeneration status of the tree species. All the study sites consist of dry deciduous forests.

2.2 Data collection and Analysis

The study was carried out in 11 study sites in each site six belt transects of 5x200 m were laid, which formed 0.6 ha in each area, and 6.6 ha in total field by study area. In each site diameter at breast height (1.37 m above ground) and height of all the individuals were recorded in each quadrat. The regeneration was studied by counting all encountered seedlings (<10 cm) and saplings (1 m height). Population structure of *C. scabiosus* was assessed by frequency histogram by categorizing the density into three gbh (girth at breast height) classes 10-20 cm, 21-30 cm and 31-40 cm. The trees were categorized into three height classes, 2-3 m, 4-5 m and 6-8 m. In all study sites associated trees with *C. scabiosus* were also recorded. Appropriate statistical analysis was done by using MS Excel.




Fig. 1. Location of the study area




Fig. 2. Dry Forests with high abundance of *Croton scabiosus* in Chitvel-Rapur, Guvvalacheruvu, Sundupalli and Pulivendula forests

3. RESULTS

A total of 2571 *C. scabiosus* individuals in 6.6 ha area, with a range of 61-100 individuals in 5x200 m quadrat (0.6 ha) were recorded. Out of these 226 are seedlings (<10 cm height), 1277 are Saplings (1 m height) and 1068 are trees (≥ 10 gbh). The number of individuals, seedlings and trees in different forests is shown in Table 1.

The gbh class distribution showed that 31.2% of 10-20 cm gbh class, 49.8% of 21-30 cm gbh

class and 19% of 31-40 cm gbh class (Fig. 3). 10-20 cm gbh class is represented by 333 trees, out of these highest in Palakondalu (20.72%) and lowest in Sanipaya (4.6%) trees. 21-30 cm gbh class represented by 532 trees. Palkondalu forest consists of percentage of trees (21.24%) of this group and lowest in Sundupalli (5.1%). 203 trees are in 31-40 cm gbh class with highest percentage in Sanipaya (32.51%) and lowest in Polatala (0.98%).

Table 1. Number of *C. scabiosus* in different forests and their Gbh and height class


Forest areas	Number of Individuals	Seedlings (<10 cm ht)	Saplings (1 m ht)	Trees (≥ 10 gbh)	Gbh class (Cm)			Height class (m)		
					10-20	21-30	31-40	2-3 m	4-5 m	6-8 m
Chitvel, -Rapur	208	20	58	136	33	65	35	50	54	32
Guvvalacheruvu	269	15	144	110	29	68	23	88	22	-
Idupulapaya	214	28	125	61	21	40	5	55	6	-
Nithyapooja kona	262	33	116	113	32	49	40	86	27	-
Palakondalu	419	16	217	178	69	113	-	163	15	-
Polathala	215	27	114	72	40	35	2	66	6	-
Porumamilla	204	8	130	66	29	35	7	51	15	-
Pulivendula	285	10	164	105	47	52	11	103	2	-
Sanipaya	328	37	133	158	15	48	66	96	54	8
Sundupalli	167	32	76	69	18	27	14	33	21	15
Jyothi	-	-	-	-	-	-	-	-	-	-
Total	2571	226	1277	1068	333	532	203	791	222	55

Table 2. Associated tree species with *C. scabiosus*

Sl. no	Tree species	Family
1	<i>Acacia caesia</i> (L.) Willd.	Mimosaceae
2	<i>Acacia chundra</i> (Rottler) Willd.	Mimosaceae
3	<i>Aegle marmelos</i> (L.) Correa	Rutaceae
4	<i>Albizia amara</i> (Roxb.) B.Boivin	Mimosaceae
5	<i>Anogeissus latifolia</i> (Roxb. ex DC.) Wall. ex Guill. & Perr.	Combretaceae
6	<i>Atalantia monophylla</i> DC.	Rutaceae
7	<i>Bauhinia racemosa</i> Lam.	Fabaceae
8	<i>Boswellia ovalifoliolata</i> N.P.Balakar. & A.N.Henry	Burseraceae
9	<i>Boswellia serrata</i> Roxb. ex Colebr.	Burseraceae
10	<i>Bridelia montana</i> (Roxb.) Willd.	Euphorbiaceae
11	<i>Buchanania axillaris</i> (Desr.) Ramam.	Anacardiaceae
12	<i>Cassia fistula</i> L.	Caesalpiaceae
13	<i>Cassine glauca</i> (Rottb.) Kuntze	Celastraceae
14	<i>Ceiba pentandra</i> (L.) Gaertn.	Bombacaceae
15	<i>Chloroxylon swietiana</i> DC.	Rutaceae
16	<i>Cleistanthus collinus</i> (Roxb.) Benth.	Euphorbiaceae
17	<i>Cochlospermum religiosum</i> (L.) Alston	Cochlospermaceae
18	<i>Commiphora caudata</i> (Wight&Arn.) Engl.	Burseraceae
19	<i>Dalbergia lanceolaria</i> L.f.	Fabaceae
20	<i>Dalbergia latifolia</i> Roxb.	Fabaceae
21	<i>Deccania pubescens</i> (Roth) Tirveng.	Rubiaceae
22	<i>Diospyros chloroxylon</i> Roxb.	Ebenaceae
23	<i>Diospyros melanoxylon</i> Roxb.	Ebenaceae
24	<i>Dolichandrone atrovirens</i> (Roth) K.Schum.	Bignoniaceae
25	<i>Erythroxylum monogynum</i> Roxb.	Erythroxylaceae
26	<i>Ficus benghalensis</i> L.	Moraceae
27	<i>Ficus mollis</i> Vahl	Moraceae
28	<i>Ficus religiosa</i> L.	Moraceae
29	<i>Ficus virens</i> Aiton	Moraceae
30	<i>Gardenia gummifera</i> L.f.	Rubiaceae
31	<i>Gardenia resinifera</i> Roth	Rubiaceae
32	<i>Grewia flavescens</i> Juss.	Tiliaceae
33	<i>Grewia hirsuta</i> Vahl	Tiliaceae
34	<i>Grewia tiliifolia</i> Vahl	Tiliaceae
35	<i>Hardwickia binata</i> Roxb.	Caesalpiaceae
36	<i>Lannea coromandelica</i> (Houtt.) Merr.	Rutaceae
37	<i>Mimosa intsia</i> L.	Mimosaceae
38	<i>Madhuca longifolia</i> (J.Kong ex L.) J.F.Macbr.	Sapotaceae
39	<i>Naringi crenulata</i> (Roxb.) Nicolson	Rutaceae
40	<i>Ochna obtusata</i> DC.	Ochnaceae
41	<i>Phyllanthus emblica</i> L.	Euphorbiaceae
42	<i>Polyalthia cerasoides</i> (Roxb.) Bedd.	Annonaceae
43	<i>Premna latifolia</i> Roxb.	Verbenaceae
44	<i>Premna tomentosa</i> Willd.	Verbenaceae
45	<i>Pterocarpus santalinus</i> L.f.	Fabaceae
46	<i>Radermarchera xylocarpa</i> (Roxb.)Roxb.ex K. Schum.	Bignoniaceae
47	<i>Santalum album</i> L.	Santalaceae
48	<i>Strychnos nux-vomica</i> L.	Loganiaceae
49	<i>Strychnos potatorum</i> L.	Loganiaceae
50	<i>Syzigium alternifolium</i> (Wight) Walp	Myrtaceae
51	<i>Terminalia bellirica</i> (Gaertn.) Roxb.	Combretaceae
52	<i>Terminalia chebula</i> Retz.	Combretaceae
53	<i>Terminalia tomentosa</i> Wight&Arn.	Combretaceae
54	<i>Wrightia tinctoria</i> R.Br.	Apocynaceae

Table 3. Associated trees with *C. scabiosus* in different forests of Kadapa districts


Forest area	Elevation (in m)	Associated Trees
Chitvel - Rapur	261-510m	<i>Limonia acidissima, Ficus virens, Mimosa intsia</i>
Idupalapaya	275m-598m	<i>Anogeissus latifolia, Dolichandrone atrovirens, Grewia hirsuta, Gardenia resinifera, Hardwickia binata</i>
Guvvalachervu	467m-519m	<i>Chloroxylon swietenia, Gardenia gummifera, Gardenia resinifera, Pterocarpus santalinus, Ochna obtusata</i>
Nithyapooja kona	407m-597m	<i>Boswellia serrata, Cleistanthus collinus, Cochlospermum religiosum, Pterocarpus santalinus.</i>
Palakondalu	155m-328m	<i>Cleistanthus collinus, Gardenia gummifera, Gardenia latifolia, Pterocarpus santalinus, Mimosa intsia, Ochna obtusata, Buchanania axillaris.</i>
Polathala	215m-368m	<i>Acacia chundra, Bridelia montana, Diospyros chloroxylon, Dolichandrone atrovirens.</i>
Porumamilla	478m-592m	<i>Cassia fistula, Diospyros melanoxylon, Phyllanthus emblica .</i>
Kanam Palli	268m-462m	<i>Anogeissus latifolia, Gardenia gummifera, Wrightia tinctoria etc.</i>
Sanipaya	205m-706m	<i>Anogeissus latifolia, Aegle marmelos, Hardwickia binata .</i>
Sundupalli	208m-515m	<i>Anogeissus latifolia, Acacia chundra, Bauhinia racemosa .</i>
Joythi	392m-635m	-

**Fig. 3. Number of stems in *C. scabiosus***

The height class distribution revealed that 74.06% trees occur in 2-3 m height class while 20.78% trees 4-5 m height class and only 5.14% trees are in 6-8 m height class (Fig. 4). Trees of 2-3 m height class are highest in Palakondalu (20.60%) and lowest in Sundupalli (4.17%). Trees of 4-5 m height class, highest in Chitvel – Rapur and Sanipaya (24.32%), lowest in Pulivendula (0.90%). Chitvel –Rapur forest is dominated by 6-8m height class (58.18%).

A total of 54 associated tree species were inventoried that belongs to 39 genera and 21 families (Table 2). Fabaceae is the dominant family represented by six species, Moraceae family with four species and Rubiaceae with

three species are the other dominant families. The other families are represented by two or one species in different forest areas. *C. scabiosus* in different forest areas commonly associated with *Acacia chundra, Boswellia serrata, Cassia fistula, Dalbergia lanceolaria, D. latifolia, Diospyros chloroxylon, D. melanoxylon, Ficus mollis, F. virens, Gardenia gummifera, G. resinifera, G. latifolia, Grewia hirsuta, G. tiliifolia, Ochna obtusa, Premna tomentosa, P. latifolia, Pterocarpus santalinus, Strychnos nux-vomica, S. potatorum* and *Wrightia tinctoria* (Table 3).

**Fig. 4. *C. scabiosus* tree height**

Over all regeneration of *C. scabiosus* is average in all the sites studied. The frequency distribution

analysis among the quadrats reveals that seedlings are present in only 35 quadrats (53.03%) out of all the 11 localities studied. The regeneration distribution shows that the saplings are 49.7% and seedlings are 8.8%. Highest regeneration status with regard to seedlings (16.37%) is observed in Sanipaya forest and lowest in Porumamilla (3.54%) and Pulivendula (4.4%) forests (Fig. 5). With regard to saplings highest percentage (17%) is in Palakondalu forest and lowest in 4.54% forests in Chitvel – Rapur (Fig. 6).

among the 11 forests varied. The size class distribution of tree has often been used to represent to the population structure of forests [11]. The forest was dominated by mostly saplings in all the 11 sites. Individuals of 10-20 cm gbh class, 21-30 cm gbh class and 2-3 m height class are more in Palakondalu (21.4%) because soil and climatic conditions are favorable for good growth of trees, and low in Sundupalli may be due to rocky habitat. Highest percentage of 31-40 cm gbh class is in Sanipaya (27.2%) this may be due good forest soil and favorable climatic conditions. This forest is undisturbed, so trees showed good growth. Lowest in Polathala (0.8%) forest which is a sacred grove with much anthropogenic disturbances. Out of total 1068 trees more

4. DISCUSSION

Population structure of *C. scabiosus* in terms of proportion of seedlings, saplings and adults




Fig. 5. Number of seedlings in different forest areas




Fig. 6. Number of saplings in different forest areas

individuals of 4-5 m height class and 6-8 m are recorded in Chitvel - Rapur hills (32.5%) low in Pulivendula (1.2%). The total basal area value is 7.74 m²/ha with a range of 25% - 75% and the average height is 3 m. *C. scabiosus* tree height and the calculated basal area showed a marked strong relationship. The total basal area recorded in the present study is 7.74 m²/ha⁻¹ and this is within the range of tropical dry forests in other parts of Eastern Ghats 6.6-23.2 m² ha⁻¹, 7.792-492.2 m² h⁻¹ [12,13].

The regeneration can be predicted by the structure of their population [14]. Though the number of population of saplings is good the seedling growth is poor. In general regeneration of species is affected by various anthropogenic activities [15,16] and natural factors such as climatic conditions. The biotic pressures play an important role in forest community dynamics and often regulate the recruitment and survival pattern of seedlings [17].

Highest percentage of seedlings growth occurred in Sanipaya (16.37%). It may be due to good soil and climatic condition favorable for growth and almost no human interference and the lowest seedling growth on rocky habitats of Pulivendula ghats (4.42%) and Porumamilla (3.53%) as the hills are rocky and lack of adequate soil for germination of seeds. Though tree population is high in Palakondalu, seedling growth in low (7.07%) in this forest. This may be due to recurrent forest fires and anthropogenic disturbances. In all areas seed germination is also affected by fruit and seed infestation and poor viability of seeds. The *C. scabiosus* tree usually occurs in lower canopy and the upper canopy is dominated by *Pterocarpus santalinus*, *Terminalia alata*, *Anogeissus latifolia*, *Gardineia gummifera*, *G. resinifera*, *Cleistanthus collinus*, *Mimosa intsia* and *Wrightia tinctoria*. *C. scabiosus* formed the dominant lower canopy tree in the elevation range of 155-328m. In Joythi forests which form the bordering area of Kadapa and Kurnool district, *C. scabiosus* trees are not found. This is in conformation with the report of Raju and Pullaiah [18].

5. CONCLUSION

The overall population structure of tree species in the study site reveals that contribution of saplings to the population was highest followed by trees and seedlings. The regeneration status of *Croton scabiosus* in various forests of Kadapa district is low. The establishment, survival and growth of

seedling is governed by several biotic and abiotic factors. Recurring forest fires and nutrient poor rocky habitat are the main abiotic factors which cause poor regeneration. Flower and fruit predation by beetles and fungal infestation of fruit and seeds, poor germination and anthropogenic disturbances are the important biotic factors for the recruitment and population expansion. Hence proper conservation measures have to be formulated to protect the species, taking into account its medicinal and economic values. Awareness programmes regarding the importance of forests, controlling forest fires and motivating the village people towards sustainable harvesting of forest products may help minimize the pressure on this species.

ACKNOWLEDGEMENTS

The authors are grateful to University Grants Commission New Delhi, India. for the financial [42-448/2013(SR)] support to the project and Forest Department, Andhra Pradesh for giving permission to work in the forests of Kadapa District.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. MoEF. National Policy and Macro level Action strategy on Biodiversity Ministry of Environment and Forest, Government of India, New Delhi; 1999.
2. Kumar N, Rita JI, Kumar N, Bhoi RK, Sajish PR. Tree species diversity and soil nutrient status in three sites of tropical dry deciduous forests of Western India. Trop. Ecol. 2010;51:273-279.
3. Khurana E, Singh JS. Germination and seedling growth of five tree species from tropical dry forests in relation to water stress: Impact of seed size. J. Trop. Ecol. 2004;20:385 -396.
4. Veenendaal EM, Swaine MO, Agyeman VK, Blay D, Abebrese IK, Mullins CE. Difference in plant and soil water relations in and around a forest gap in West Africa during the dry season may influence seedling establishment and survival. J. Ecol. 1996;83:83-90.
5. Ahmedullah M, Nayar MP. Endemic plants of the Indian Region. Peninsular India, Botanical survey of India, Calcutta. 1987;1.

6. Babu, PSP. Euphorbiaceae. In: Pullaiah T, Moulali DA. Flora of Andhra Pradesh, India. Vol. 2. Scientific Publishers, Jodhpur. 1997;836-890.
7. Salamma S, Rao BRP. Distribution and Conservation status of *Croton scabiosus* Bedd. (Euphorbiaceae) an endemic tree species of South Eastern Ghats of Andhra Pradesh, India. J. Threatened Taxa. 2014; 6(10):6363-6370.
8. Balakrishnan NP, Chakrabarthym T. The family of Euphorbiaceae in India. Bishen Singh Mahendra Pal Singh, Dehradun; 2007.
9. Devi SN, Mani J, Venkata Raju RR, In Vitro antimicrobial activity and phytochemical constituents of *Croton scabiosus* Bedd. Intern. J. Pharmaceut. Res. Dev. 2011;3(6):149-154.
10. Neginhal SG. Forest trees of South India. Navabharath press, Bangalore; 2014.
11. Saxena AK, Singh JS. Tree Population structure of certain Himalayan forest association and implications concerning their future composition. Vegetation. 1984; 58(2):61-69.
12. Jha CS, Singh JS. Composition and dynamics of dry tropical forest in relation to soil. J. Vegetation Sci. 1990;1:609-614.
13. Reddy CS, Parchi Ugle. Tree species diversity and distribution pattern in tropical forest of Eastern Ghats, India; case study. Life sciences Journal. 2008;4:87-93.
14. Khan ML. Population structure of some tree species in disturbed and protected sub tropical forests of North east India. Acta Oecologia Applicata. 1987;8(3):247-255.
15. Barik SK. Dynamics of tree seedling population in a humid sub tropical forest of North east India as related to disturbances, condition. J. Forest Research. 1986;26(4): 584-589.
16. Iqbal K. Regeneration status of tree species around Khoh river in Garhwal Himalaya, Indian J. Forestry. 2012;35(4): 471-476.
17. Pala NA. Diversity and regeneration status of Sarkot Van panchyat in Garhwal Himalaya, India. J. Forestry Res. 2012; 23(3):399-404.
18. Raju RRV, Pullaiah T. Flora of Kurnool (Andhra Pradesh). Bishen Singh Mahendra Pal Singh, Dehradun; 1995.

© 2017 Nagireddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/18693>