



# Effect of Edible Coating on Shelf-life Enhancement of Papaya (*Carica papaya* L.) at Ambient Conditions

Ritu Bakhla <sup>a\*</sup>, Annjoe V. Joseph <sup>b\*\*</sup> and Samir Topno <sup>b\*\*</sup>

<sup>a</sup> Horticulture (Fruit Science), Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, (UP), India.

<sup>b</sup> Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, (UP), India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i143062

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100938>

Original Research Article

Received: 27/03/2023  
Accepted: 30/05/2023  
Published: 07/06/2023

## ABSTRACT

An experiment was conducted on the effect of edible coating on shelf-life enhancement of papaya (*Carica papaya* L.) at ambient conditions. The main aim of this study was to increase the shelf life of papaya by using edible coating. Postharvest losses in fruits is a serious problem due to rapid deterioration during storage. Use of edible coatings over fruits is used to improve their quality and enhance the shelf life. The main purpose of this study was to assess the effect of aloe vera gel, beeswax, paraffin wax and gelatine coating in extending the shelf-life of papaya including physico-chemical changes under ambient storage conditions. The experiment was laid out in completely randomized design (CRD) with 7 treatments and 3 replications. The treatment consisted of coating

\*\* Assistant Professor;

\*Corresponding author: E-mail: rbakhla3@gmail.com;

material with Aloe vera gel 0.5% (T1), Aloe vera gel 1%(T2), Aloe vera gel 1.5%(T3), Beeswax 6% (T4), Paraffin wax 100% (T5), Gelatin10%(T6), Control(T7) and freshly harvested papaya fruits. Statistical analysis revealed that among all the treatments, the maximum effect on extending the shelf-life and delaying the ripening was recorded in paraffin wax coated papaya fruits. Therefore, coating of papaya fruits with paraffin wax can be suggested as a post-harvest treatment to enhance the shelf-life.

**Keywords:** Papaya; edible coatings; physico-chemical characteristics; shelf-life.

## 1. INTRODUCTION

Papaya (*Carica papaya* L.) is a tropical fruit belonging to the family Caricaceae. Papaya is a nutritious table fruit of high digestive value and rich in vitamins and minerals. Papaya fruit are rich in enzymes called papain and chymopapain. Both enzymes digest protein, meaning they can help with digestion and reduce inflammation. Papaya is a highly nutritious and rich source of Vitamin A, B1, B2, E and C, minerals like potassium, magnesium, boron, iron, calcium, and phosphorus, and copper. Further, it also contains 85-90% water, 10-13% sugar and 0.6% protein. The energy value of papaya is 200 kJ/100 g. Sugars present in papaya include glucose, fructose and sucrose. Papaya contains high levels of antioxidants that reduce the risk of heart diseases. The antioxidants prevent the oxidation of cholesterol. Papaya has folic acid which is essential for converting the amino acid homocysteine into less harmful amino acid.

Papaya is categorized as a climacteric fruit with an increased respiration rate during ripening process and its respiration rate can be inhibited by providing a coating on the surface of the fruit including aloe vera gel, beeswax, paraffin wax, gelatine etc.

“Due to the high perishable nature of papaya, a major portion of the production goes into waste which is reported to be around 40-60% of the total production at various papaya growing regions” [1]. Increasing shelf life minimizes the loss of water, oxygen, and other soluble material from the fruits and vegetables. Edible coated fruits can have several advantages such as retention of colour, TSS, acidity, flavour for a longer period, reduction of post-harvest disorders; disease, therefore, increases the shelf life of the product.

“Aloe vera (*Aloe barbadensis* Miller) is a perennial plant of the Liliaceae family with turgid green leaves joined at the stem in a rosette pattern. Aloe vera gel is tasteless, colourless and

odourless. This natural product is a safe and environmentally friendly alternative to synthetic preservatives such as sulphur dioxide” [2].

“Beeswax is derived from honeycomb and is produced by honey extraction. A study by Purwoko and Fitriadesi [3] showed that beeswax coating at 6% concentration can reduce weight loss and total soluble solids of papaya Solo cv Tainung compared to control after 14 days of storage”.

“Paraffin wax is a distillate of crude petroleum [4] and is used for coating raw fruits and vegetables. Paraffin wax is used to make fruits, vegetables and candy look shiny. Food-grade paraffin wax is composed of vegetable oils, palm oil derivatives, synthetic resins and other materials”.

“Gelatin, an important biopolymer obtained by hydrolysis from collagen, is widely used with a broad range of functional properties and applications such as in the food, pharmaceutical and photographic industries, including its film-forming ability. Gelatin films generally have effective barrier properties against oxygen and carbon dioxide. Aguilar-Mendez et al. [5] found that the application of gelatin-starch coatings extended the postharvest shelf life of avocado”.

The main purpose of this study was to assess the effect of aloe vera gel, beeswax, paraffin wax and gelatine coating in extending the shelf-life of papaya including physico-chemical changes at ambient storage conditions. The study also aimed at analysing the physical and chemical changes in papaya during storage at ambient conditions.

## 2. MATERIALS AND METHODS

The experiment on enhancing the shelf life of the papaya was conducted at Postharvest Laboratory, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, and PRAYAGRAJ (UP) during 2022-2023.

The experimental design was CRD (Completely Randomized Design) with 7 treatments and 3 replications. Freshly harvested papaya fruit of variety the red lady with uniform fruit shape and size was used. The coated fruits were kept at ambient condition for 15 days and observation on physiological loss in weight, total soluble solid, pH, vitamin C/ascorbic acid, titratable acidity, fruit colour, fruit texture, fruit taste, fruit appearance, fruit shelf life and fruit overall acceptability was taken at 3 days interval. Observations on fruit colour, fruit texture, fruit taste, fruit appearance and overall acceptability were evaluated from hedonic rating test with a panel of 5 members to make judgment more accurate.

The data recorded during the course of investigation were subjected to statistical analysis using CRD (Completely Randomized Design) as per the method of "Analysis of Variance" technique. The significant difference between the treatment's means were tested at 5% level of significance.

The overall significance of differences among the treatments was tested, using critical difference (C. D. at 5%) level of significance. The result was statistically analysed with the help of a window based computed package OPSTAT (Sheoran, 2004).

### 3. RESULTS AND DISCUSSION

#### 3.1 Effect of Edible Coating on Physiological Loss of Weight during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied.

The data is illustrated in Table 1, which indicates that the treatment T5 (Paraffin wax) showed lowest physiological loss in weight (4.64%, 6.22%, 7.26%, 8.30% and 10.30%) at 3<sup>rd</sup>, 6<sup>th</sup>, 9<sup>th</sup>, 12<sup>th</sup> and 15<sup>th</sup> days of storage followed by Beeswax coating. Similarly, T7(control) indicates the highest physiological loss in weight of (7.51%, 8.33%, 10.22%, 12.18%, 13.86%).

"Increase in PLW with the storage period might be due to loss of moisture from the fruits by way of transpiration or evaporation. The paraffin wax serves as a physical barrier around the fruit which partially closes the stomatal openings and lenticels thereby reducing the rates of

transpiration and respiration" [6]. "The highest rate of PLW in control might be due to higher moisture loss humidity as compared to coated fruits" [7]. Similar findings have also been reported by Hazarik et al. [8].

#### 3.2 Effect of Edible Coating on Total Soluble Solid during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied.

The data is illustrated in Table 2, which indicates that the treatment T5 (Paraffin wax) showed the highest TSS of (9.99<sup>0</sup>brix) and lowest TSS was indicated in T7 control of (6.72<sup>0</sup>brix). "Total soluble solid of the papaya increased with the increase in storage period from 3<sup>rd</sup> day to 12<sup>th</sup> day of storage and thereafter it decreased indicating rapid metabolic breakdown in those fruits. The increase in TSS during storage may possibly be due to breakdown of complex organic metabolites into simple molecules or due to hydrolysis of starch into sugars. On complete hydrolysis of starch, no further increase in sugars occurs and subsequently a decline in these parameters is predictable as they along with other organic acids are the primary substrate for respiration" [9]. "The increase in TSS in wax coated fruits till 12th day and then gradual decrease as compared to control fruits indicates the possible role in delaying metabolic activities of fruits during ripening and storage" [10]. Similar findings have also been reported by Suwanti et al. [11].

#### 3.3 Effect of Edible Coating on pH during Storage in paPaya

The statistical analysis revealed significant differences among the different treatments studied.

The data is illustrated in Table 3, which indicates that the T7 (Control) showed the highest pH value of (6.45) and the lowest pH value was indicated in T5 (6.11).

The pH of papaya increases during storage. An increase in pH papaya during storage was due to the conversion process of carbohydrate to sugar and the use of acids for metabolism. Similar findings have also been reported by Suwanti et al. [11].

**Table 1. Effect of edible coating on physiological loss of weight during storage in papaya**

Treatment	Treatment details	Physiological loss of weight				
		No. of days at room temperature				
		Initial	6 Days	9 days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	6.36	7.34	9.34	10.20	12.59
T <sub>2</sub>	Aloe vera gel (1%)	6.74	7.34	9.38	11.55	12.16
T <sub>3</sub>	Aloe vera gel (1.5%)	6.59	7.39	9.22	10.36	11.58
T <sub>4</sub>	Beeswax (6%)	5.23	6.39	7.40	8.22	10.51
T <sub>5</sub>	Paraffin wax (100%)	4.64	6.22	7.26	8.30	10.30
T <sub>6</sub>	Gelatin (10%)	5.74	6.34	7.38	9.22	10.85
T <sub>7</sub>	Control	7.51	8.33	10.22	12.18	13.86
	C.D. at 5%	0.34	0.23	0.25	0.50	0.94

**Table 2. Effect of edible coating on total soluble solid during storage in papaya**

Treatment	Treatment details	Total soluble solid (° Brix)				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	6.40	7.71	9.02	10.33	7.30
T <sub>2</sub>	Aloe vera gel (1%)	5.78	7.09	8.40	9.71	7.35
T <sub>3</sub>	Aloe vera gel (1.5%)	5.46	6.77	8.11	9.35	7.35
T <sub>4</sub>	Beeswax (6%)	5.11	6.42	7.73	9.01	7.69
T <sub>5</sub>	Paraffin wax (100%)	4.75	6.06	7.37	8.61	9.99
T <sub>6</sub>	Gelatin (10%)	5.15	6.46	7.77	9.04	7.39
T <sub>7</sub>	Control	6.81	8.12	9.45	10.74	6.72
	C.D. at 5%	1.02	1.02	1.03	1.03	0.73

**Table 3. Effect of edible coating on pH during storage in papaya**

Treatment	Treatment details	pH				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	4.73	5.30	5.65	5.76	6.26
T <sub>2</sub>	Aloe vera gel (1%)	4.62	5.21	5.36	5.51	6.20
T <sub>3</sub>	Aloe vera gel (1.5%)	4.49	5.20	5.31	5.41	6.15
T <sub>4</sub>	Beeswax (6%)	4.35	5.33	5.44	5.50	6.12
T <sub>5</sub>	Paraffin wax (100%)	4.36	5.28	5.40	5.50	6.11
T <sub>6</sub>	Gelatin (10%)	4.43	5.26	5.44	5.53	6.15
T <sub>7</sub>	Control	4.77	5.58	5.73	6.10	6.45
	C.D. at 5%	0.08	0.09	0.17	0.24	0.16

**Table 4. Effect of edible coating on vitamin C/ascorbic acid(mg/100gm) during storage in papaya**

Treatment	Treatment details	Vitamin C /ascorbic acid (mg / 100g)				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	34.78	40.67	43.58	42.65	35.13
T <sub>2</sub>	Aloe vera gel (1%)	35.12	40.54	43.50	42.56	35.20
T <sub>3</sub>	Aloe vera gel (1.5%)	35.15	40.42	43.35	42.45	35.29
T <sub>4</sub>	Beeswax (6%)	33.21	41.16	42.19	45.49	36.17
T <sub>5</sub>	Paraffin wax (100%)	45.48	51.24	56.97	67.99	62.80
T <sub>6</sub>	Gelatin (10%)	35.09	40.25	43.53	42.66	35.16
T <sub>7</sub>	Control	35.32	40.66	45.71	42.82	33.02
	C.D. at 5%	0.58	0.59	0.79	0.39	0.69

### 3.4 Effect of Edible Coating on Vitamin C/ Ascorbic Acid during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied. The data is illustrated in Table 4, which indicates that the T5 (paraffin wax) showed highest retention of vitamin C of (62.80 mg/100g) and lowest retention of vitamin C was indicated in T7 control (33.02 mg/100g). Vitamin C of papaya in all treatments increased until it reaches the peak maturity “The fruits coated with paraffin wax show high retention of vitamin C due to retardation of oxidation process and consequently slow rate of conversion of L-ascorbic acid into dehydroascorbic acid by ascorbic acid oxidase. The retention of higher ascorbic acid in coated fruits might be due to the ripening retarding effect and slow rate of biological activities during storage”. [11] Similar observations have also been recorded in mango [6] and mandarin orange [12].

### 3.5 Effect of Edible Coating on Titratable Acidity during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied.

The data is illustrated in Table 5, which indicates that the T5 (paraffin wax) showed highest retention of titratable acidity of (1.70%) and lowest retention of titratable acidity was indicated in T7 control (1.45%). “The titratable acidity of the fruits decreased with the increase of storage period. The decrease in TA with the increase in storage period could possibly be attributed due to the decreased hydrolysis of organic acids in pyruvate decarboxylation reaction occurring during the ripening process and subsequent accumulation

of organic acids which were oxidised at a slow rate because of decreased respiration”. [8] Similar findings have also been reported by [13].

### 3.6 Effect of Edible Coating on Taste during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied.

The data is illustrated in Table 6, which indicates that the T5 (paraffin wax) showed highest retention of taste (7.46) which shows that it is liked moderately followed by Beeswax (7.33) and lowest retention of taste was indicated in T7 Control (5.13) which shows that it is neither liked nor disliked. “The papaya fruits showed a gradual and steady increase in the taste of the fruits up to the 12th day of storage and after that a sharp decline was observed. The paraffin wax coated fruits were rated as extremely desirable after 12 days and thereafter taste declined. The gradual increase in the taste of papaya fruits during storage has been attributed to the increase in the concentration of total sugars and TSS contributing to the typical papaya taste”. [8] Similar findings have also been reported by Hazarik et al. [8].

### 3.7 Effect of Edible Coating on Texture during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied.

The data is illustrated in Table 7, which indicates that the T5(paraffin wax) showed highest retention of texture (8.16) which shows that it is liked very much and lowest retention of texture in

**Table 5. Effect of edible coating on titratable acidity during storage in papaya**

Treatment	Treatment details	Titratable acidity (%)				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	1.17	1.29	1.43	1.66	1.51
T <sub>2</sub>	Aloe vera gel (1%)	1.25	1.38	1.49	1.68	1.59
T <sub>3</sub>	Aloe vera gel (1.5%)	1.32	1.39	1.52	1.71	1.59
T <sub>4</sub>	Beeswax (6%)	1.42	1.52	1.63	1.77	1.65
T <sub>5</sub>	Paraffin wax (100%)	1.51	1.65	1.72	1.82	1.70
T <sub>6</sub>	Gelatin (6%)	1.37	1.47	1.57	1.66	1.59
T <sub>7</sub>	Control	1.12	1.34	1.50	1.78	1.45
	C.D. at 0.5%	0.09	0.09	0.09	0.09	0.13

**Table 6. Effect of edible coating on taste of during storage in papaya**

Treatment	Treatment details	Taste				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	6.29	6.62	6.62	6.84	5.46
T <sub>2</sub>	Aloe vera gel (1%)	6.33	6.57	6.57	6.71	5.36
T <sub>3</sub>	Aloe vera gel (1,5%)	6.39	6.54	6.54	6.68	5.50
T <sub>4</sub>	Beeswax (6%)	7.68	7.79	7.79	8.49	7.33
T <sub>5</sub>	Paraffin wax (100%)	8.38	8.54	8.54	8.69	7.46
T <sub>6</sub>	Gelatin (10%)	6.36	6.53	6.53	6.64	5.50
T <sub>7</sub>	Control	5.67	6.11	6.11	6.52	5.13
	C.D. at 5%	0.16	0.19	0.19	0.15	0.27

**Table 7. Effect of edible coating on texture during storage in papaya**

Treatment	Treatment details	Texture				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	6.37	6.84	6.17	5.81	5.63
T <sub>2</sub>	Aloe vera gel (1%)	6.50	6.71	6.30	6.22	5.83
T <sub>3</sub>	Aloe vera gel (1.5%)	6.72	6.68	6.54	6.43	6.24
T <sub>4</sub>	Beeswax (6%)	7.45	8.49	7.25	7.17	7.11
T <sub>5</sub>	Paraffin wax (100%)	8.82	8.69	8.59	8.43	8.16
T <sub>6</sub>	Gelatin (10%)	6.90	6.64	6.62	6.55	6.40
T <sub>7</sub>	Control	6.82	6.52	6.48	5.86	5.69
	C.D. at 5%	0.25	0.15	0.18	0.19	0.17

T7 control (5.69) which shows that it is neither liked nor disliked. "The paraffin wax coatings of fruits resulted in higher fruit texture, which might be due to reduced transpiration and respiration activity along with delayed ethylene production and thus retained more turgidity of the cells of the fruits" [14]. Our study is in close conformity with the findings of [15] and [16].

### 3.8 Effect of Edible Coating on Colour during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied.

The data is illustrated in Table 8, which indicates that the T5 (paraffin wax) showed highest retention of colour which is (7.46) which shows that it is liked moderately followed by T4 Beeswax (7.33) and lowest retention of colour was indicated in T7 control (5.13) which shows that it is neither liked nor disliked. "In all the treatments, the colour of the fruits changed from

green to yellow from the first day of storage (3rd day) till the end of storage period (16 days). In wax coated fruits, the development of yellow colour was at a slow and steady rate indicating more consumers preferred peel colour till the end of storage period. In wax coated fruits, the development of yellow colour increased at a slower rate till 15th day of storage resulting in development of uniform coloured fruit surface". [8] Similar findings have also been reported by Hazarik et al. [8].

### 3.9 Effect of Edible Coating on Appearance during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied.

The fruits treated with paraffin wax (T5) had gloss, shiny appearance and no wrinkles and browning, therefore scoring maximum value with respect to appearance (7.67) which shows that it is liked moderately followed by T4 Beeswax

(6.68) at 15 days of storage (Table 9). On the other hand, T7 control fruits registered the minimum appearance value (5.220) which shows that it is neither liked nor disliked. The development of better appearance in the wax coated fruits could be possibly due to creation of favourable gaseous atmosphere under congenial temperature. Our study is in close conformity with the findings of Ladaniya [17] in Mosambi sweet oranges where the wax coated fruits had better appearance as compared to other fruits.

### 3.10 Effect of Edible Coating on Overall Acceptability during Storage in Papaya

The statistical analysis revealed significant differences among the different treatments studied.

The data is illustrated in Table 9, which indicates that the T5 (paraffin wax) showed the highest score for overall acceptability which is (7.51) which shows that it is liked moderately while the lowest was observed in T7 control (4.610) which shows that it is disliked slightly. The papaya fruits showed a gradual and steady decline in the overall acceptability of the fruits with the increase

in storage period. The wax coated fruits were rated as extremely desirable after the 12th day and thereafter taste declined. The overall acceptability of papaya fruits depends on a delicate balance of sugars, acids, phenolics and aromatic compounds with a number of additional factors such as pulp texture and visual appearance also influence the perceived quality and consumer acceptance and appreciation. Similar findings have also been reported by Hazarik et al. [8].

### 3.11 Effect of Edible Coating on Shelf-life during Storage in Papaya

The shelf-life of papaya fruits varied significantly among different treatments. Among all the treatments, fruits coated with paraffin wax T5 have maximum shelf-life of 15 days followed by T4 Beeswax 12 days and T3 Aloe vera gel 1.5% coatings with 9 days, while control recorded the lowest shelf-life of 6 days. The probable reason for paraffin wax coated fruits having maximum shelf-life might be due to its ability to serve as a physical barrier around the fruit which reduces transpiration and respiration activity along with delayed ethylene production.

**Table 8. Effect of edible coating on colour during storage in papaya**

Treatment	Treatment details	Colour				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	6.15	6.29	6.62	6.84	5.46
T <sub>2</sub>	Aloe vera gel (1%)	6.21	6.33	6.57	6.71	5.36
T <sub>3</sub>	Aloe vera gel (1.5%)	6.29	6.39	6.54	6.68	5.50
T <sub>4</sub>	Beeswax (6%)	7.57	7.68	7.79	8.49	7.33
T <sub>5</sub>	Paraffin wax (100%)	8.28	8.38	8.54	8.69	7.46
T <sub>6</sub>	Gelatin (10%)	6.25	6.36	6.53	6.64	5.50
T <sub>7</sub>	Control	5.40	5.67	6.11	6.52	5.13
	C.D. at 5%	0.13	0.16	0.19	0.15	0.27

**Table 9. Effect of edible coating on appearance during storage in papaya**

Treatment	Treatment details	Appearance				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	6.36	6.24	6.19	6.08	5.67
T <sub>2</sub>	Aloe vera gel (1%)	6.50	6.39	6.29	6.19	6.08
T <sub>3</sub>	Aloe vera gel (1.5%)	6.71	6.61	6.54	6.39	6.15
T <sub>4</sub>	Beeswax (6%)	7.45	7.32	7.22	7.16	6.68
T <sub>5</sub>	Paraffin wax (100%)	8.53	8.46	8.36	8.10	7.67
T <sub>6</sub>	Gelatin (10%)	6.96	6.86	6.77	6.60	6.49
T <sub>7</sub>	Control	6.83	5.63	5.49	5.40	5.22
	C.D. at 5%	0.28	0.27	0.26	0.26	0.20

**Table 10. Effect of edible coating on overall acceptability during storage in papaya**

Treatment	Treatment details	Overall acceptability				
		No. of days at room temperature				
		Initial	6 Days	9 Days	12 Days	15 Days
T <sub>1</sub>	Aloe vera gel (0.5%)	6.38	6.25	5.66	5.50	4.95
T <sub>2</sub>	Aloe vera gel (1%)	6.81	6.65	6.38	5.66	5.35
T <sub>3</sub>	Aloe vera gel (1.5%)	6.84	6.69	6.49	6.25	5.71
T <sub>4</sub>	Beeswax (6%)	7.32	7.20	7.08	6.69	6.46
T <sub>5</sub>	Paraffin wax (100%)	8.37	8.25	8.12	7.68	7.51
T <sub>6</sub>	Gelatin (10%)	6.88	6.71	6.57	6.39	6.25
T <sub>7</sub>	Control	6.30	5.70	5.65	5.33	4.61
	C.D. at 5%	0.04	0.12	0.24	0.204	0.29

“In addition, waxing filled the lenticels and left no places for entry of pathogens like fungi and bacteria so ultimately it increases the shelf life of the papaya fruits. The positive effect of wax coating on storage life could probably be due to the modification of the atmosphere. The modified atmosphere created could, therefore, delay the ripening by delaying ethylene production and by reducing the level of internal oxygen and consequently prolonging the storage life of papaya fruit” Gol and Rao et al. [18] [19,20].

#### 4. CONCLUSION

Among all the tested treatments, paraffin wax has minimum PLW (10.30%), with high TSS value of (9.99<sup>o</sup>brix), with maximum vitamin C content (62.80mg/100gm) and maximum titratable acidity of (1.70%) and the lowest pH value was indicated in T5 (6.11). And in organoleptic parameters the highest score for taste (7.46) which is liked moderately, texture (8.16) which is liked very much, colour (7.46) which is liked moderately, appearance (7.67) which is liked moderately and overall acceptability (7.51) which is liked moderately was also recorded in T5 (Paraffin wax). Among all the tested treatments, Paraffin wax has enhanced storage life capacity up to 15 days and retained the nutritional quality as compared to the fruit of the control set. The maintenance of quality and extension of shelf life of papaya fruit by paraffin wax demonstrate that such a coating can be considered for commercial application during its storage.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Prasad K, Paul J. Postharvest losses of papaya and practice for its management. 2021;2:49–53.
2. Serrano M, Valverde JM, Guillén F, Castillo S, Martínez-Romero D, Valero D. Use of aloe vera gel coating preserves the functional properties of table grapes. *Journal of Agriculture and Food Chemistry*. 2006;54(11):3882–3886.
3. Purwoko BS, Fitriadesi P. Effect of Coating materials and storage temperatures on quality and shelf life of papaya fruit. *Buletin Agronomi*. 2000;28:66-72.
4. Bennett H. *Industrial Waxes*. New York: Chemical Publishing company; 1975.
5. Aguilar-Mendez MA, Martin-Martinez ES, Tomas SA, Cruz-Orea A, Jaime-Fonseca MR. Gelatin-starch films: Physicochemical properties and their application in extending the post-harvest shelf life of avocado (*Persea americana*), *Journal of the Science of Food and Agriculture*. 2008;88:185–193.
6. Jain Shashi, Mukherjee S. Enhancing keeping quality of fruits in mango cv. Langra. *Indian Journal of Horticulture*. 2011;68:142-144.
7. Pongener Alemwati, Mahajan BVC, Singh Harminder. Effect of different packaging films on storage life and quality of peach fruits under cold storage conditions. *Indian Journal of Horticulture*. 2011;68:240-245.
8. Hazarik TK, Lalthanpuui D, Mandal. Influence of edible coatings on physico-chemical characteristics and shelf-life of papaya (*Carica papaya*) fruits during ambient storage. *Indian Journal of Agricultural Sciences*. 2017;87(8): 1077–83.
9. Wills RBH, Bembridge PA, Scott KJ. Use of flesh firmness and other objective tests



- to determine consumer acceptability of delicious apples. Australian Journal of Experimental Agriculture and Animal Husbandry. 1980;20:252–6.
10. Fan X, Blankenship SM, Mattheis JP. 1-MCP inhibits apple ripening. Journal of American Society for Horticultural Science. 1999;124:690–5.
  11. Suwanti R Utami, Kawiji Praseptianga, Khasanah. The effect of packaging methods (paper, active paper, and edible coating) on the characteristic of papaya MJ9 ambient temperature storage. IOP Conference Series: Earth and Environment al Science. 2018;102:012087.
  12. Yadav M, Kumar N, Singh DB, Singh GK. Effect of post-harvest treatments on shelf-life and quality of Kinnow mandarin. Indian Journal of Horticulture. 2010;67:243–8.
  13. Pinal B Vyas, Neeta B Gol, Ramana Rao TV. Postharvest quality maintenance of papaya fruit using polysaccharide-based edible coatings. International Journal of Fruit Science. 2014;14:1:81-94.
  14. Nanda S, Rao DVS, Krishnamurthy S. Effect of shrink film wrapping and storage temperature on the shelf life and quality of pomegranate fruits cv Ganesh. Postharvest Biology and Technology. 2001;22:61–9.
  15. Da Luz, Lucas, Vettorazzi, Júlio, Santa, Renato, & Barros, Fabio, Barros, Gislanne, Pereira, Messias, Cardoso, Deisy. Sensory acceptance and qualitative analysis of fruits in papaya hybrids. Anais da Academia Brasileira de Ciências. 2018; 90:10:1590
  16. Chaturvedi Devendra Kumar, Shukla Rama. Evaluation of qualitative attributes of papaya leather. International Journal of Agricultural Engineering. 2018;11: 84-89.
  17. Ladaniya MS. Response of 'Mosambi' sweet orange (*Citrus sinensis*) to degreening, mechanical waxing, packaging and ambient storage conditions. Indian Journal of Agricultural Sciences. 2001;71:234–9.
  18. Gol NB, Rao TVR. Banana fruit ripening as influenced by edible coatings. International Journal of Fruit Science. 2011;11:119–35.
  19. Sarkar HN, Hasan MA, Chattopadhyay PK. Effect of packaging and chemicals on the storage behavior of sapota. Horticulture Journal. 1995;8:17–24.
  20. Sindhu GS, Dhillon WS, Mahajan BVC. Effect of waxing and packaging on storage of pear cv. Punjab Beauty. Indian Journal of Horticulture. 2009;66:239–44.

© 2023 Bakhla et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/100938>